



National Institute for Public Health
and the Environment
Ministry of Health, Welfare and Sport

Physical damage due to drug dependence

RIVM report 340041001/2012

J.G.C. van Amsterdam | E.J.M. Pennings |

T.M. Brunt | W. van den Brink



National Institute for Public Health
and the Environment
Ministry of Health, Welfare and Sport

Physical damage due to drug dependence

RIVM Report 340041001/2012

Colophon

© RIVM 2012

Parts of this publication may be reproduced, provided acknowledgement is given to the 'National Institute for Public Health and the Environment', along with the title and year of publication.

J.G.C. van Amsterdam, RIVM, Laboratory for Health Protection
Research

E.J.M. Pennings, The Maastricht Forensic Institute

T.M. Brunt, Trimbos Institute

W. van den Brink, Academisch Medisch Centre - Amsterdam Institute
for Addiction Research

Contact:

Dr. J.G.C. van Amsterdam

RIVM-GBO

jan.van.amsterdam@rivm.nl

This investigation has been performed by order and for the account of ZonMw (Netherlands Organisation for Health Research and Development) within the framework of ZonMw programme 'Risk behavior and Dependence'.

Abstract

Physical damage due to drug dependence

Excessive use of alcohol, tobacco and illicit drugs cause long term physical health damage. Smoking can cause lung cancer, and alcohol abuse may lead to liver cirrhosis and tumors in the oral cavity, esophagus and liver. The medical treatment of the diseases leads to financial costs. The health effects and treatment of alcohol and tobacco use are higher as compared to those of illicit drugs use. This is mainly due to the relatively large number of people smoking (27 percent of the Dutch population) or excessive alcohol use (84 percent drink, of which 10 percent drink excessive). Illicit drugs are, however, used by relatively few people (recent use is 0.1 to 4.2 percent) and mostly used for only some years. At the individual level, the physical health effects of alcohol and tobacco use are similar to those of recreational use of hard drugs heroin and crack. In general, only the intensive use of illicit drugs i.e. the abuse of drugs leads to great physical health damage.

Physical effects of drugs are limited

This emerged from a literature review of the RIVM, which gives an overview of the physical health damage of seventeen recreational drugs, alcohol and tobacco. The study was commissioned by ZonMw, that has also initiated research to the psychological and social effects of illicit drugs. The latter consequences are often greater than the physical health damage.

Physical effects of the four most commonly used drugs

Of the four most commonly used illicit drugs, like ecstasy, cocaine, cannabis, amphetamine, ecstasy seems not to lead to serious physical health damage. Cannabis smoking can cause lung cancer and COPD. The use of heroin, cocaine and crack can cause infectious diseases, AIDS and tuberculosis; the use of dirty needles caused the most problems here. Cocaine, crack and (repeated) amphetamine use is related to heart disease. Of all illicit drugs, risk of (fatal) heart attack is greatest when snorting cocaine. The use of khat and anabolic steroids is associated with cardiovascular disease, but the evidence is fairly weak. Urological complications by regular use of the anesthetic ketamine are reported in the literature, but they are rare. Oral cancers can be caused by intensive use khat. Finally, almost all the problematic hard drug users suffer from dental disease.

For most illicit drugs, it is difficult to indicate the association between the use and the diseases caused by them, because the drugs are frequently combined with other drugs, tobacco and alcohol (poly drug use). In particular, the extent to which and how the illicit drugs are used in the past, is hardly known. This knowledge is necessary to link disease to the use of various drugs.

Keywords:

illicit drugs, physical effects, disease, relative risk, alcohol, tobacco

Rapport in het kort

Lichamelijke gevolgen van druggebruik

Overmatig gebruik van alcohol, tabak en drugs veroorzaakt op termijn lichamelijke gezondheidsschade. Roken verhoogt het risico op longkanker en overmatig alcoholgebruik is geassocieerd met levercirrose en tumoren in de mondholte, slokdarm en de lever. De medische behandeling daarvan brengt kosten met zich mee. Ten opzichte van druggebruik zijn de gezondheidsschade en de behandelingskosten van overmatig alcohol- en tabakgebruik hoger. Dat komt vooral doordat relatief veel mensen roken (27 procent van de Nederlandse bevolking) of overmatig alcohol gebruiken (84 procent drinkt, waarvan 10 procent overmatig). Drugs worden daarentegen door relatief weinig mensen (recent gebruik is 0,1 tot 4,2 procent) en meestal gedurende enkele jaren gebruikt. Op individueel niveau is de lichamelijke gezondheidsschade van overmatig alcohol- en tabakgebruik vergelijkbaar met die van recreatief gebruik van de harddrugs heroïne en crack. In het algemeen leidt alleen intensief gebruik van drugs en genotmiddelen tot grote lichamelijke gezondheidsschade.

Lichamelijke gevolgen van drugs zijn beperkt

Deze conclusie blijkt uit een literatuuronderzoek van het RIVM. Hierin wordt een overzicht gegeven van de lichamelijke gezondheidsschade van zeventien recreatieve drugs, alcohol en tabak. Het onderzoek is uitgevoerd in opdracht van ZonMw, dat ook onderzoek heeft laten doen naar de psychische, verslavende en sociale effecten van drugs. Deze gevolgen zijn waarschijnlijk vaak groter dan de lichamelijke gezondheidsschade.

Lichamelijke gevolgen van de vier meest gebruikte drugs

Van de vier meest gebruikte drugs, cocaïne, cannabis, amfetamine en ecstasy, lijkt het gebruik van ecstasy niet te leiden tot ernstige lichamelijke gezondheidsschade. Het roken van cannabis is positief geassocieerd met longkanker en COPD. Het gebruik van heroïne, cocaïne en crack kan leiden tot infectieziekten, AIDS en tuberculose; het gebruik van vuile naalden veroorzaakt hierbij de meeste problemen. Cocaïne-, crack- en (herhaald) amfetaminegebruik is gerelateerd aan hart- en vaatziekten. Van alle drugs is kans op een (fatale) hartaanval het grootst bij het snuiven van cocaïne. Ook het gebruik van khat en anabole steroïden wordt in verband gebracht met hart- en vaatziekten, maar het bewijs hiervoor is vrij zwak. Urologische complicaties door regelmatig gebruik van het narcosemiddel ketamine worden in de literatuur gemeld, maar ze komen weinig voor. Orale kankervormen kunnen ontstaan door intensief khatgebruik. Tot slot hebben vrijwel alle problematische gebruikers van harddrug tandheelkundige aandoeningen.

Voor de meeste drugs is het moeilijk om aan te geven wat het verband is tussen het gebruik en de ziekten die daaruit voortvloeien. De drugs worden namelijk vaak in combinatie met andere drugs, en met tabak en alcohol gebruikt (polydruggebruik). Vooral de mate waarin en wijze waarop de middelen in het verleden zijn gebruikt, is nauwelijks bekend. Deze kennis is nodig om een verband te kunnen leggen tussen ziekten en het gebruik van de verschillende genotmiddelen.

Trefwoorden:

drugs, lichamelijke effecten, ziekte, relative risico, alcohol, tabak

Contents

1	Samenvatting—9
2	Kennishiaten—11
3	Conclusion—13
4	Gaps in knowledge—15
5	General introduction—17
5.1	Prevalence of drug use in The Netherlands—17
5.2	Polydrug use—17
5.3	Infectious disease—18
5.4	Cardiovascular disease due to drugs abuse—18
5.5	Hyperthermia due to drug abuse in general—18
5.6	Periodontal disease—19
5.6.1	Characteristics of the patient group—19
5.6.2	Determinants of periodontal disease—19
5.6.3	Prevalence of periodontal disease—19
5.6.4	Practical problems—20
5.7	Other addictions—20
5.8	Financial burden of drug addiction—21
6	Methylphenidaat (Ritalin®)—23
6.1	Acute adverse effects—23
6.2	Chronic adverse effects—23
6.3	Diseases—24
7	Anabolic steroids (AAS)—25
7.1	Acute adverse effects—25
7.2	Chronic adverse effects—25
7.3	Disease—26
7.3.1	Cancer disease—26
7.3.2	Liver disease—26
7.3.3	Cardiovascular disease—26
7.3.4	Other diseases—27
8	Benzodiazepines—29
8.1	Acute adverse effects—29
8.2	Chronic adverse effects—29
8.3	Disease—29
9	Khat—31
9.1	Acute adverse effects—31
9.2	Chronic adverse effects—31
9.2.1	Overview—31
9.2.2	Cardiovascular complications—31
9.2.3	Oral and gastro-intestinal complications—32
9.2.4	Growth retardation—32
9.3	Diseases—32
9.3.1	Cancer disease—32
9.3.2	Periodontal disease—33

- 9.3.3 Reproductive effects—33
- 10 LSD—35**
 - 10.1 Acute adverse effects—35
 - 10.2 Chronic adverse effects—35
 - 10.3 Diseases—35
- 11 Magic mushrooms—37**
 - 11.1 Acute adverse effects—37
 - 11.2 Chronic adverse effects—37
 - 11.3 Diseases—37
- 12 Ketamine—39**
 - 12.1 Acute adverse effects—39
 - 12.2 Chronic adverse effects—39
 - 12.3 Diseases—39
- 13 GHB (Gamma Hydroxy Butyric acid)—41**
 - 13.1 Acute adverse effects—41
 - 13.2 Chronic adverse effects—41
 - 13.3 Diseases—41
- 14 Ecstasy—43**
 - 14.1 Acute adverse effects—43
 - 14.2 Chronic adverse effects—43
 - 14.3 Diseases—43
- 15 Cannabis—45**
 - 15.1 Acute adverse effects—45
 - 15.2 Chronic adverse effects—45
 - 15.2.1 Pulmonary system—45
 - 15.2.2 Immune System—45
 - 15.2.3 Endocrine System—45
 - 15.3 Disease—45
 - 15.3.1 Pulmonary disease—45
 - 15.3.2 Cancer disease—46
 - 15.3.3 Cardiovascular disease—46
 - 15.3.4 Reproductive disorders—47
 - 15.3.5 Other diseases—47
 - 15.4 Health benefits—47
- 16 Amphetamine—49**
 - 16.1 General remarks on amphetamine and methamphetamine—49
 - 16.2 Acute adverse effects—49
 - 16.3 Chronic adverse effects—50
 - 16.3.1 Cardiovascular pathology—50
 - 16.4 Disease—50
 - 16.4.1 Cardiovascular disease—50
 - 16.4.2 Infections—51
 - 16.4.3 Other diseases—51
- 17 Methamphetamine—53**
 - 17.1 Acute adverse effects—53
 - 17.2 Disease—53
 - 17.2.1 Cardiovascular disease—53

- 17.2.2 Periodontal disease—53
- 17.2.3 Dermatological disorders—54

18 Opiates—55

19 Buprenorphine—57

- 19.1 Acute adverse effects—57
- 19.2 Chronic adverse effects—57
- 19.3 Diseases—57

20 Methadone—59

- 20.1 Acute adverse effects—59
- 20.2 Chronic adverse effects—59

21 Heroin—61

- 21.1 Effects of short-term use—61
 - 21.1.1 Use at lower doses—61
 - 21.1.2 Use at higher doses—61
- 21.2 Chronic adverse effects—61
- 21.3 Disease following chronic heroin use—62
 - 21.3.1 Pulmonary disease—62
 - 21.3.2 Infections—62
 - 21.3.3 Renal disease—63
 - 21.3.4 Other diseases—63
- 21.4 Mortality—63
- 21.5 Economical burden—63

22 Cocaine—65

- 22.1 Acute adverse effects—65
- 22.2 Chronic adverse effects—65

23 Cocaine-crack—67

- 23.1 Acute adverse effects—67
- 23.2 Chronic adverse effects—67
- 23.3 Disease following chronic cocaine and crack use—67
 - 23.3.1 Cardiovascular diseases—68
 - 23.3.2 Pulmonary disease—68
 - 23.3.3 Kidney disease—69
 - 23.3.4 Infections related to needle use—69
- 23.4 Mortality—69
- 23.5 Economic burden—69

24 Alcohol—71

- 24.1 Acute adverse effects—71
- 24.2 Disease due to chronic excessive alcohol use—71
 - 24.2.1 Introduction—71
- 24.3 General protective and harmful effects—72
 - 24.3.1 Disease in the digestive tract (non-carcinogenic)—73
 - 24.3.2 Cardiovascular disease—74
 - 24.3.3 Cancer disease—75
 - 24.3.4 Hematologic/hematopoietic system—76
 - 24.3.5 Other diseases—76
- 24.4 Mortality—76
- 24.5 Financial burden related to alcohol consumption—76

25	Tobacco—79
25.1	Acute adverse effects—79
25.2	Diseases—79
25.2.1	Cancer disease—79
25.2.2	Respiratory Disease—80
25.2.3	Cardiovascular disease—80
25.2.4	Other diseases—80
25.3	Environmental tobacco smoke—80
25.4	Mortality—81
25.5	Economic burden—81
26	Acknowledgements—83
27	Accountability—85
28	References—87

1 Samenvatting

Voor de meeste recreatieve drugs is het moeilijk om sterke associaties te vinden tussen het gebruik van de drugs en de mogelijke ziekte die daaruit voortvloeit. De belangrijkste reden is dat veel recreatieve drugs in combinatie met andere drugs worden gebruikt (polydrugsgebruik), zodat de associaties zwakker worden. Een duidelijker beeld ontstaat voor de drugs die relatief vaak gebruikt worden, zoals heroïne, cocaïne, cannabis, amfetamine en ecstasy. Van deze vijf drugs lijkt het gebruik van cannabis en ecstasy niet echt te leiden tot heftige lichamelijke aandoeningen. Een uitzondering is het roken van cannabis, dat aan kanker van de onderste en bovenste luchtwegen, en COPD gerelateerd is. Vermoedelijk zijn stoffen in de tabaksrook, anders dan de cannabinoïden, verantwoordelijk voor deze carcinogene effecten.

Vaak voorkomende ziektes na parenteraal gebruik van heroïne, cocaïne en crack zijn infectieziekten, zoals AIDS en tuberculose. Deze ziekten worden niet door de psychoactieve stoffen veroorzaakt, maar door de vuile naalden (als gevolg van het uitwisselen van naalden), die bij het gebruik van de drugs gebruikt worden. Het gebruik van cocaïne en crack is specifiek gerelateerd aan cardiotoxiciteit, wat leidt tot hart- en vaatziekten, hartinfarct, ritmestoornissen en cerebrovasculair accident (CVA). Ook (herhaald) amfetaminegebruik kan leiden tot hart- en vaatziekten.

Het recreatieve gebruik van methylfenidaat, LSD, benzodiazepines, paddo's en GHB leidt niet tot lichamelijke aandoeningen, hoewel sommige van deze drugs wél geassocieerd zijn met psychische stoornissen. Het gebruik van khat en anabole steroïden wordt in verband gebracht met hart- en vaatziekten, maar het bewijs hiervoor is vrij zwak. De orale kankervormen die door khatgebruik kunnen ontstaan zijn waarschijnlijk te wijten aan verbindingen in de khat anders dan de psychoactieve componenten. Ecstasy kan vrijwel zonder lichamelijke consequenties worden gebruikt, mits de gebruiker regelmatig water of frisdrank drinkt (met mate; niet overmatig) om oververhitting te voorkomen. Urologische complicaties door regelmatig ketaminegebruik worden in de literatuur gemeld, maar de incidentie is laag. Bovendien is de relatie onduidelijk, omdat ketamine vaak in combinatie met andere drugs gebruikt wordt (polydrugsgebruik). Tot slot lijden vrijwel alle problematische hard drug gebruikers tandheelkundige aandoeningen.

De conclusie van het onderzoek is dat recreatief druggebruik nauwelijks leiden tot lichamelijke ziektes. In Nederland worden zij zelden gemeld, hetgeen een aanwijzing is dat het Nederlandse drugsbeleid gericht op 'harm reduction' nog steeds zeer effectief is. Bij heftig gebruik (vaak gebruik en/of in hoge doseringen) kan het gebruik van de drugs wél 'problematisch' en leiden tot soms ernstige lichamelijke ziektes.

Betrouwbare gegevens over de financiële lasten (en de DALY's) van recreatief druggebruik zijn niet beschikbaar. Bovendien kan door het vaak voorkomende polydruggebruik het oorzakelijke verband tussen ziekte en het gebruik van een bepaald geneesmiddel niet worden gegeven. Ten tweede is het drugsgebruik van de patiënt in de jaren die vooraf gingen aan de openbaring van de ziekte grotendeels onbekend.

Sommige schattingen van de financiële last van overmatig alcoholgebruik en tabaksgebruik tonen aan dat de medische kosten van deze twee en de meest

gebruikte 'drugs' relatief hoog zijn ten opzichte van de last voortkomend uit het gebruik van illegale drugs.

2 Kennishiaten

Een panel van experts heeft, mede op geleide van dit rapport, de belangrijkste kennishiaten in de kennis over lichamelijke effecten van drugs geïdentificeerd en geprioriteerd. Deze zijn 'Welke interventies zijn bewezen als zijnde effectief', 'De toxiciteit van polydrugsgebruik', 'Lange-termijn effecten van coma-zuipen' en 'De mogelijke neurotoxiciteit van GHB'.

Prioriteits-score *	Hiaat in het kort	Beschrijving van de kennishiaat
3.8	Effectieve interventies	Beschrijf de 'hoog risico groepen', de bewezen en effectieve interventies voor deze groepen (in wie effectief; in wie niet) en hoe hoog zijn de kostenbesparingen van deze interventies? Zijn het per definitie de 'vroege interventies'?
3.7	Toxiciteit van polydrug gebruik	Beschrijf de toxiciteit en de ziektes tengevolge van polydruggebruik met in het bijzonder de combinatie van drug(s) met alcohol?
3.5	'Coma-zuipen'	Wat zijn de lange termijn effecten (op de hersenen, cognitie) van herhaalde coma's als gevolg van overmatig alcoholgebruik ('binge drinken') door jongeren?
3.4	GHB neurotoxiciteit	Wat zijn de schadelijke effecten op langere termijn van herhaaldelijk in coma geraken tengevolge van GHB-overdoseringen?
3.2	Problematisch druggebruik	Hoe hoog is de prevalentie van problematisch drugsgebruik? Wat is het effect van een selectieve reductie op middelen, ziektelast en de behandelingskosten? Modelmatige benadering
3.0	Kosten en DALY's	Hoe hoog zijn de medische en maatschappelijke kosten (DALY's) van (elk van) de recreatieve drugs (waaronder populaire combinaties bij poly-druggebruik)?
2.4	Cardiotoxiciteit van cocaïne	Breng de cardiotoxiciteit van cocaïne in kaart, met inbegrip van leefstijl en type gebruik. Mogelijk is het gebruik van cocaïne geassocieerd met honderden fatale hartaanvallen in Nederland (per jaar is 40% van alle fatale hartaanvallen bij mannen van 25-40 jaar gerelateerd aan cocaïnemisbruik).
2.3	Mondhygiëne	Mondhygiëne is een belangrijk onderdeel van de kwaliteit van leven van druggebruikers. Er zijn geen gegevens over in welke mate een goede mondhygiëne hun kwaliteit van leven en terugkeer in de samenleving kan verbeteren.
2.2	Prevalentie gegevens	Hoe hoog is de prevalentie van de minder vaak gebruikte drugs (LSD, khat, methyfenidaat, anabole steroïden)?
1.9	Alternatieve doseringen	Hoe hoog is de haalbaarheid van minder schadelijke alternatieve toedieningsvormen (ontwikkeling, onderwijs en kosten-effectiviteit)?

* Na het bepalen van tien thema's door de veertien experts gaven de experts voor elk thema een score op een schaal van 0 (geen prioriteit) tot 5 (hoogste prioriteit). Voor meerdere thema's kon de hoogste score worden gegeven.

3 Conclusion

Regarding the low prevalence of various diseases in relation to recreational drug use in The Netherlands, physical disease burden in of drug related disease is relatively low. It thus appears that the Dutch 'harm reduction' drug policy has been and still is very effective.

Solid data about the financial burden (and DALY's) of recreational drug use are not available. In addition, due to the highly prevalent polydrug use, the causal relation between disease and the use of a specific drug cannot be given.

Secondly, the history of drug use by the patient is largely unknown. Some estimates of the financial burden of alcohol over-consumption and tobacco use show that the medical costs of treatment are relatively high for these two most prevalently used 'drugs'.

The major gaps in knowledge identified are 'Which are the proven effective interventions', 'Toxicity of polydrug use', 'Long-term effects of binge drinking ("coma-zuipen")' and 'Potential neurotoxicity of GHB'.

4 Gaps in knowledge

Forteen experts were invited to define gaps in knowledge and to subsequently score the gaps. The panel of experts had to cover the wide variety of items characterising the harm profile of the different drugs reviewed. Therefore experts with the following disciplines were recruited from the network of the authors: oncology-intensive care, internist/toxicology, intensivist, drug addiction care, pharmacology, epidemiology and sociology working in the field of illicit drugs. The four most prominent gaps in knowledge are 'Which interventions have proved to be effective', 'Toxicity of polydrug use', 'Long-term effects of comatose excessive alcohol consumption' and 'Possible neurotoxicity of GHB'. The following ten gaps in knowledge have been identified:

Priority score *	Gap in short	Description of the Gap
3.8	Effective interventions	Describe the groups at high risk and what are the proven and effective interventions for these groups (in which group effective; in which group ineffective) and what are the reductions in costs by these interventions? Are they by definition 'early interventions'?
3.7	Toxicity of polydrug use	Describe the toxicity and disease due to polydrug use, more specifically the combination of drug(s) with alcohol?
3.5	Binge drinking ("Coma-zuipen")	What are the long-term effects (on the brain and cognition) of repeated coma's as a result of excessive alcohol consumption (binge drinking) by young people?
3.4	GHB neurotoxicity	What are the long-term adverse effects of repeatedly slipping in coma as a result of GHB overdosing?
3.2	Problematic use of drugs	What is the prevalence of problematic drug use? What would be the effect of selective reduction on resources, disease burden and treatment costs. Model-based approach.
3.0	Costs and DALY's	What are the medical and societal costs (DALY's) of (each of) the recreational drugs (including popular combinations in poly-drug use)?
2.4	Cardiotoxicity of cocaine	Map the cardiotoxicity of cocaine, including contributing factors (lifestyle, type of use). Possibly, cocaine use is associated with hundreds of fatal heart attacks in The Netherlands (per annum 40% of all fatal heart attacks in men 25-40 years is linked to cocaine abuse).
2.3	Oral health	Oral health is an important aspect of the quality of life of drug users. Data are lacking on how this issue improves their quality of life and rehabilitation in society.
2.2	Prevalence data	What is the prevalence of the less frequently used drugs (LSD, khat, methylphenidaat, anabolic steroids)?
1.9	Alternative dosages	What is the feasibility of alternative dosage forms, which are less harmful? Development, education and cost efficiency?

* Experts (N=14) could score the ten items on a scale from 0 (no priority) to 5 (top priority); a top score could be given to more than one item.

5 General introduction

Illicit drugs, alcohol and tobacco provide significant physical co-morbidity and mortality worldwide. The literature on the associations of substance use and physical (or somatic) medical illness is large, but often limited to the main illegal drugs. Some of the reviews looked at these co-morbidity or co-occurring disorders, but no comprehensive and systematic review on this topic is currently available.

This present review describes state of the art of physical disorders related to recreational drug abuse. Mental, psychiatric and social burden of drug abuse are addressed in separate state-of-the-art studies currently prepared by others. Clinical signs of overdosing the recreational drugs and damage to the unborn child are no topic of this review.

5.1 Prevalence of drug use in The Netherlands

The prevalence of drug use in The Netherlands in 2010 according National Drug Monitor (NDM, 2011).

Drug	Prevalence 2010 (%)			Remarks
	Life time	Last year	Last month	
Cannabis	25.7	7.0	4.2	(almost) daily use by 30%
Cocaine	5.2	1.2	0.5	
Heroin (and methadon)	0.5	0.1	0.1	
Ecstasy	6.1	1.4	0.4	
Amphetamine	3.1	0.4	0.2	
GHB	1.3	0.4	0.2	
Alcohol		84	76	10% is heavy drinker *
Tobacco	60	-	27.1	6.3% is heavy smoker

* 32% is binge drinker (last six months prevalence)

Prevalence of other drugs abuse, like methylfenidaat (Ritalin), buprenorphine and benzodiazepines is not known. Crack-cocaine and ketamine are rarely used (the latter 'only' by 'psychonauts' and clubbers). Methamphetamine is also rarely used and limited to a few scenes (homosexuals, psychonauts). Dutch figures are not available. Magic mushrooms: ever use 3% (2002), last month 0.3% (2001), but a sharp decrease since the drug was banned. LSD ever use is 1.4%, and last month 0.1% (2005). Khat is used only by African-Arab immigrants (lifetime 78%; last month 34-67%). The lifetime prevalence of anabolic steroids reported is 0%-6%. Some 50,000 sports players use anabolic steroids regularly.

5.2 Polydrug use

Many cocaine and amphetamine abusers (about 60% to 80%) simultaneously drink alcohol (Heil et al., 2001, Pennings et al., 2002). People who are opioid dependent tend towards alcohol when their primary drug is not available because alcohol may boost the effects of other drugs (Coffin et al., 2003, Kandel and Davies, 1996). Probably as many as half of the men and a quarter of the women with opioid dependence became also dependent on alcohol within the first five years after active opioid involvement. GHB users often also use

ketamine. Ecstasy (and amphetamine) is often used in combination with alcohol or downers to dampen the ecstasy (and amphetamine) effects. Another example is AAS (anabolic steroid) abuse. It appears that the adjusted odds ratio (95% C.I.) for reported 12-month marijuana use by college students was 2.9 (1.7, 5.0) (McCabe et al., 2007). For other drugs the odds ratios ranged from 6.1 (ecstasy) to 13.0 (cocaine). AAS users were also more likely to be alcohol dependent with an odd ratio of 3.1 (1.7, 5.7). Many other examples and reasons for polydrug use can be given, but this is beyond the scope of this review.

5.3 Infectious disease

The association of recreational drugs use and the increased incidence of infections (hepatitis viruses, human immunodeficiency virus and sexually transmitted pathogens) are known, but it is less clear whether the current drug of abuse itself is the causal factor. It is very likely that the intravenous route of administration of illegal drugs (cocaine, heroin, methamphetamine) is the co-morbid link, not the drug itself. For details see the specific sections.

5.4 Cardiovascular disease due to drugs abuse

Drug-induced hypertension is an important risk factor of stimulant drug abuse, because it can cause stroke (usually cerebral haemorrhage), acute myocardial infarction, pulmonary oedema, dissecting aneurysm, and/or hypertensive encephalopathy (Eagle et al., 2002, Qureshi et al., 1988a). Among patients with stroke due to cocaine abuse about 50% have cerebral haemorrhage, 30% subarachnoid haemorrhage, and 20% have ischemic stroke (Mueller et al., 1990, Tardiff et al., 1989). In San Francisco General Hospital, drug abuse was identified as the most common predisposing condition among young patients (<35 years of age) presenting with stroke (Kaku and Lowenstein, 1990c). Most of the patients had either infective endocarditis (13/73) or stroke occurring soon after the use of a stimulant (34/73). The relative risk for stroke among drug abusers, adjusted for other stroke risk factors, was estimated as 6.5 (Kaku and Lowenstein, 1990a). Vasculitis has been associated with nearly every drug of abuse (Citron et al., 1970, Rumbaugh et al., 1971, Rumbaugh et al., 1976), including heroin (Brust and Richter, 1976, Woods and Strewler, 1972), methylphenidate (Trugman, 1988) and cocaine (Fredericks et al., 1991, Kaye and Fainstat, 1987, Krendel et al., 1990, Treadwell and Robinson, 2007).

5.5 Hyperthermia due to drug abuse in general

It does not take long either to boil an egg or to cook neurons (Hamilton, 1976). Severe hyperthermia (>40.5°C) is generally recognized as cause of major morbidity and mortality, regardless of the cause. Various drugs (Callaway and Clark, 1994, Ebadi et al., 1990, Lecci et al., 1991, Sporer, 1995, Walter et al., 1996) can cause hyperthermia, and this may initially be overlooked while the more familiar manifestations (i.e., seizures) of the intoxication are being managed. Classic heat stroke is characterized by a body temperature of $\geq 40.5^{\circ}\text{C}$, and severe CNS dysfunction has been associated with a chance of disabling neurologic sequelae and with mortality rates of up to 80% (Sarnquist and Larson, 1973).

All amphetamines including amphetamine, methamphetamine, and MDMA can produce lethal hyperthermia (Gordon et al., 1991, Jaehne et al., 2005). Large overdoses of LSD have also been associated with severe hyperthermia (Friedman and Hirsch, 1971, Klock et al., 1974b).

5.6 Periodontal disease

5.6.1 *Characteristics of the patient group*

In the Netherlands, it is estimated that there are 350,000 problem drinkers, 185,000 heavy drinkers, and 20,000 to 30,000 hard drug users. There are approximately 25,000 primary heroin addicts, of which some 13,000 participate in a methadone programme (Hendriks et al., 2003) and nearly three-fourth has regular contact with social workers (Goppel et al., 2003).

The Centre for Special Dentistry of Jellinek (CBT) is the only specialised dental clinic for heavy drug users in The Netherlands. Around 3,500 patients are treated annually in the CBT. Note that this group is the 'top of the iceberg', as one may assume that all hard drug users have similar dental problems, but do - for various reasons - not use this clinic. Of addicts seeking help at the CBT, 90% was polydrug user, 83% used heroin and/or cocaine, 50% used intravenously, 68% used methadone, 23% used amphetamines and 16% used hallucinogens. Only 10% was addicted to alcohol. Of injecting users, 90% had hepatitis B and/or hepatitis C, 13% suffered from endocarditis, and 20% is HIV and/or AIDS positive. The psychopathology of most patients in the CBT treatment Jellinek is very high: 95% of the patients meet the DSM-IV criteria for some mental disorder. Heroin users and alcoholics often suffer from mental depression and addiction-related stress, while cocaine users are often manic and so often restless, irritable and hyper-alert.

5.6.2 *Determinants of periodontal disease*

Of the patients visiting the CBT for the first time, 37% has pain (Ter Horst et al., 1999), whereas in a regular practice this is only 2% (Den Dekker, 1990). Beside the 37%, some 18% gives cavities as a reason to visit the CBT (Molendijk et al., 1995c, Ter Horst et al., 1999).

Poor oral health among drug addicts can be seen as a reflection of various adverse factors such as malnutrition, inadequate oral hygiene, and consuming many sweets. Drug addicts consume large amounts of sugar and have no more regular meals (Carter, 1978). Most (84%) of American heroin addicts used in their addiction more sugar (Picozzi et al., 1972). In a Dutch study, 37% of the heroin addicts answered that they used sugar more than 15 times a day (Molendijk et al., 1995b).

Psychotropic drugs have a dampening pain effect. As the pain damping effect disappears during and following abstinence, the drug user becomes aware - via a toothache - of the bad dental condition. The severe sensation can lead to a relapse, because the patient misses the relief of the drugs with analgesic properties.

5.6.3 *Prevalence of periodontal disease*

According to a Dutch study, 81% of alcoholics and drug addicts have much plaque (more than 1/3 of the buccal or lingual tooth surface) (Molendijk et al., 1995a), which may be explained by the combination of poor oral hygienic behaviour, a sugar-rich diet and decreased salivary secretion (Scheutz, 1984). Addicts have a relative high caries prevalence as shown by the higher DMFS (Decayed, Missing, or Filled Surface) score of 52.1 as compared to the general population (score of 38.9) (Molendijk et al., 1995d). The high score in the addicts was mainly due to the increased number of carious surfaces (sixfold) and extracted teeth or elements (1.8-fold) (Molendijk et al., 1995e). This difference was no longer significant when addicts are compared with groups with low socioeconomic status, which implies that drug dependence is only one of the

determinants of dental decay. Alcoholics show a higher caries prevalence as compared to the general population, as well (Friedlander and Gorelick, 1988a). The use of stimulants, such as alcohol and cocaine, leads to bruxation habits with attrition of teeth (Friedlander and Gorelick, 1988b, Lee et al., 1991a). Together with reduced function by missing teeth, this can result in temporo-mandibular dysfunction. The use of cocaine, especially in combination with alcohol, also can induce severe xerostomia (dry mouth) (Lee et al., 1991b). Local application of cocaine can cause vestibular and gingival damage, erosion or ulceration (Parry et al., 1996). Smoking crack can cause the same picture, especially on the palate (Mitchell-Lewis et al., 1994).

When cocaine addicts are under the influence of the drug, they tend - in their mania - to scrub their teeth so violently that the elements abrade and cervical and gingival lacerations are produced. Excessive alcohol consumption leads to chronic gastritis with increased acid production and reflux, and frequent vomiting, which causes erosion of the teeth.

A recent prospective study in New Zealand (Thomson et al., 2008) showed that subjects with the highest exposure to marijuana (N=182) had the highest number of incident attachment losses, and the highest incidence of one or more attachment loss (CAL) sites ≥ 4 mm (RR 1.6; 95% CI: 1.2-2.2) compared with a no-exposure group (N=293) (controlled for tobacco smoking, irregular use of dental services, and dental plaque).

Reduced loss of aesthetic elements is a major mental problem for the patient, especially when the elements of the front teeth are missing. Understandably, the patient prefers to fill the diastema, but the priorities of severely mutilated teeth lie elsewhere at that time. Obviously, the refurbishment of the teeth can contribute to a positive self-image of the patient (Molendijk, 1992). In this way, the dental treatment has a positive impact on the rehabilitation process of the addict.

Finally, it should be noted that the impact of poor oral health in drug addicts on quality of life is much higher as compared to that of subjects where either an impacted wisdom tooth has been surgically extracted or of those with severe gingivitis (van Wijk et al., 2011).

5.6.4 *Practical problems*

The treatment of addicts is not limited to dental problems, because the dentist also engages specific drug related medical complications and psycho-social problems, including antisocial personality disorders (maladaptive behaviours, aggression of the patient) and anxiety (especially intra-oral injections). 40 per cent of the patients do not turn up at an appointment (33). The main reasons are: (1) long-term abuse of alcohol (and ketamine) which can result in memory loss, (2) psychopathology, (3) fear of dental treatment, (4) no currently available assistance for visits/transport to the CBT, or (5) because of other priorities, such as need to score drugs (Sainsbury, 1999).

5.7 **Other addictions**

In addition to illicit drugs (chemicals), subject may be dependent on certain 'habits or hobbies', like gambling, internet, food over-consumption, intensive sporting, bulimia. High speed car driving is not regarded as addiction, but rather as a bad habit.

It is quite obvious that food related disorders may lead to obesity or extreme low body weight. Intensive sporting (e.g. marathon) may be sound for the pulmonary condition, but certainly not for the locomotor apparatus (knees, feet). Gambling and internet addiction may have a profound impact on mental

health, but not on physical health. It may lead to poverty leading to a poor nutrition and physical health.

5.8 Financial burden of drug addiction

For most drugs, no clear figures about drug related costs (hospital care, general health care) are available. Alcohol and tobacco are an exception here. The main reason is polydrug use, so that the direct association between the use of a certain drug and the disease a drug user suffers from remains unclear. In addition, the prevalence of use of most drugs is too low to attain sufficient power in the statistical analysis.

6 Methylphenidaat (Ritalin®)

Methylphenidate is a safe drug when used in the recommended oral dose. Methylphenidate is structurally related to amphetamine; as compared to amphetamine it stimulates the central nervous system less potently and with more mental than motor effects, and has minimal peripheral effects in therapeutic doses.

6.1 Acute adverse effects

In children, Ritalin induces loss of appetite and sleep disturbances. The loss of appetite may be so severe that growth is significantly impeded, but both these side effects usually disappear after reducing the dose and ensuring that the first dose of the day is given after rather than before breakfast.

In a large dose, it stimulates the central nervous system and elicits convulsions. It is more potent than amphetamine as an antidepressant, and in exacerbating schizophrenic symptoms. Occasionally, anorexia, nausea, dry mouth, nervousness, insomnia, dizziness, and palpitation have been recorded (Iversen, 2008).

Methylphenidate, like amphetamines and amphetamine-like drugs which act on the peripheral sympathetic nervous system, increases the heart rate and blood pressure. Normally this is hardly relevant, but there have been reports of serious adverse events associated with the cardiovascular system and even some deaths. Cardiac dysrhythmias, shock, cardiac muscle pathology, and liver pathology have all been reported (Chernoff et al., 1962). When given parenterally, methylphenidate may increase blood pressure and/or pulse rate more frequently (Witton, 1964). Occasionally, methylphenidate causes abdominal distress, which can be reduced by lowering the dose or by administration immediately after meals (Greenhill et al., 2002, Lopez et al., 2003). As with other stimulants, chorea (Extein, 1978) and choreoathetosis (Weiner et al., 1978) can be precipitated in children and adults at higher methylphenidate doses. Hypersensitivity reactions have been reported. Skin reactions have included exfoliative dermatitis and erythema multiforme. Purpura, thrombocytopenia, and leucopenia have occurred. Blood counts should be monitored periodically during prolonged therapy.

6.2 Chronic adverse effects

Little information exists about the long-term effects of methylphenidate. Methylphenidate has been reported to cause stunting of growth by impairing growth hormone secretion (Holtkamp et al., 2002). One study showed that methylphenidate produced decreases in weight percentiles after 1 year of therapy and progressive decrement in height percentiles that became significant after 2 years of use (Mattes and Gittelman, 1983). However, another study suggested that moderate doses might have a lower risk for long-term height suppression than dexamphetamine (Greenhill et al., 1984). Though methylphenidate retarded growth rate during active treatment, final height was not compromised and that a compensatory rebound of growth appeared to occur on stopping stimulant treatment (Klein and Mannuzza, 1988), confirming that there is no evidence for long-term growth impairment (Hechtman and Greenfield, 2003).

6.3 Diseases

Stimulants like methylphenidate may be associated with cardiac complications (Jaffe and Kimmel, 2006c, Lucas et al., 1986). Of 289 patients exposed to excessive doses of methylphenidate, none of the patients developed severe symptoms, although 31% showed symptoms, like tachycardia, and agitation (White and Yadao, 2000). Crushed tablet preparations meant for oral use, especially methylphenidate (Ritalin) to inject the drug give rise to foreign body emboli in the circulation and lodge in the lung, forming granulomas. Granulomas may also form in the lung and brain (possibly due to the passage of foreign materials through a patent foramen ovale) which may require surgical intervention.

7 Anabolic steroids (AAS)

Steroid abuse disrupts the normal production of hormones in the body, causing both reversible and irreversible changes. Side effects of AAS, however, develop virtually only during long term use (Thiblin and Petersson, 2005). The most common side-effects are cosmetic in nature and reversible. Class B AAS cause hepatic toxicity (Welder et al., 1995) leading to jaundice after two to five months, but hepatotoxicity has never been described with the parenteral use of testosterone esters. Severe side effects on the liver and lipoproteins mainly result from alkylated AAS at high dose (Ishak and Zimmerman, 1987), whereas parenteral AAS appear to damage heart muscles which may become clinically prominent after several years. Most of the serious life-threatening effects appear relatively infrequent.

7.1 Acute adverse effects

Minor acute side effects of steroid use are: head aches, fluid retention (especially in the extremities), gastrointestinal irritation, diarrhoea, stomach pains, premature male pattern baldness and an oily skin. Acute effects with some more clinical impact are jaundice, menstrual abnormalities, and hypertension. Infections can develop at the injection site, causing pain and abscess. In both sexes, acne develops at puberty (i.e. not in adults) during treatment with androgens due to the growth of sebaceous glands and the secretion of the natural oil sebum (Karila et al., 2003b). Observational studies suggest that a majority (88-96%) of anabolic steroid users experience at least one objective side-effect, including acne (40-54%), testicular atrophy (40-51%), gynaecomastia (10-34%), cutaneous striae (34%) and injection-site pain (36%) (Evans, 2004). Usually, HPT function recovers within weeks to months, but persisting hypogonadism (for more than a year after AAS discontinuation) has been described (Boyadjiev et al., 2000, Menon, 2003a, Van Breda et al., 2003).

7.2 Chronic adverse effects

Health consequences associated with anabolic steroid abuse include urogenital problems, acne, and cardiovascular and hepatic disease (Melchert and Welder, 1995, Rogol and Yesalis, 1992, Sullivan et al., 1998a). AAS supplements suppress the hypothalamic-pituitary-testicular (HPT) axis in males, so that discontinuation – especially abruptly after a prolonged use – can cause them to become hypogonadal. Males using high doses of AAS can have the circulating estrogen levels typical of women during a normal menstrual cycle (Wilson, 1988a), which can lead to breast pain in men and the often irreversible gynaecomastia. Gynaecomastia, particularly when painful, may require surgical correction. Changes in males that can be reversed include reduced sperm production, impotence, difficulty or pain in urinating and shrinking of the testicles (testicular atrophy). In one study of male bodybuilders, more than half had testicular atrophy and/or either reversible or irreversible breast development (gynaecomastia) (Wilson, 1988b). In females, elevated AAS levels result in menstrual irregularities and the development of more masculine characteristics such as decreased body fat and breast size, deepening of the voice, excessive growth of body hair (such as moustache and beard growth), and irreversible loss of scalp hair (baldness), as well as clitoral enlargement. With continued administration of steroids, clitoral hypertrophy and deepened voice become irreversible (Shifren, 2004, Wilson, 1992).

7.3 Disease

Medical complaints as described under or resulting from chronic toxicity regularly occur, and are experienced as very unpleasant and disturbing. Their frequency depends on the dose and the length of the period of use.

7.3.1 *Cancer disease*

Anabolic steroid use has been associated with prostate cancer (Creagh et al., 1988). Of particular concern is premature physeal closure in any child/adolescent, which results in a decrease in adult height. In some cases, however, AAS is clinically used to limit the abnormal body length. AAS give an increased risk for fatal liver cysts, other liver changes, and liver cancer (Bagia et al., 2000, Gorayski et al., 2008, Kafrouni et al., 2007, Sanchez-Osorio et al., 2008, Socas et al., 2005, Velazquez and Alter, 2004). AAS effects on the prostate effects include hypertrophy (Jin et al., 1996, Wemyss-Holden et al., 1994) and perhaps an increased risk of prostate cancer (Anonymous, 1991, Roberts and Essenhig, 1986), although the latter association has been questioned (Morgentaler, 2006, Morgentaler, 2007).

7.3.2 *Liver disease*

Class B and C AAS are highly hepatotoxic. The alkylated AAS have also been shown to increase hepatic triglyceride lipase activity between 21% and 123% and low density lipoprotein by as much as 29% (Bagatell and Bremner, 1996a, Thompson et al., 1989c). On the other hand, AAS-induced hepatic pathology is often reversible upon discontinuation of AAS (Modlinski and Fields, 2006), and the overall prevalence of adverse hepatic effects among long-term AAS users is likely low (Pope and Katz, 1994).

7.3.3 *Cardiovascular disease*

The long-term use of AAS has been reported to be associated with cardiovascular disease (CVD), like hypertension, heart attack and stroke, but the prevalence and underlying mechanisms of AAS-induced cardiovascular toxicity remain poorly understood. Steroids contribute to the development of CVD, partly by changing the levels of lipoproteins that carry cholesterol in the blood. Steroids, particularly oral steroids, increase the level of low-density lipoprotein cholesterol (LDL-cholesterol) and decrease the level of high-density lipoprotein cholesterol (HDL-cholesterol). Notably the alkylated and orally used AAS such as stanozolol (6 mg per day p.o. for six weeks) lower HDL-cholesterol by 33%, particularly HDL2-cholesterol which is reduced 23 to 80% (Bagatell and Bremner, 1996b, Thompson et al., 1989b). The effect of parenteral testosterone enanthate (200 mg per week for six weeks) itself is much less dramatic, with only a 9% reduction in HDL-cholesterol (Thompson et al., 1989a). In general, serum levels return to baseline level within several weeks to months after drug cessation (Hartgens and Kuipers, 2004). Even the administration of high doses of testosterone enanthate (600 mg per week) parenterally for 20 weeks hardly affected HDL (Singh et al., 2002). The increase in cholesterol is likely to be associated with narrowing of the arteries, i.e. atherosclerosis, and subsequent heart attacks.

Indeed, power lifters have a greater risk of atherosclerosis secondary to increased concentrations of LDL-cholesterol and decreased concentration of HDL-cholesterol (Hurley et al., 1984). In addition, steroids induce blood clotting due to increased platelet count and aggregation (Ferenchick et al., 1992, Togna et al., 2003). Steroids can also cause myocardial hypertrophy, which also increases the likelihood of arrhythmias, sudden death, systolic and diastolic

hypertension, and myocardial infarct (Frankle et al., 1988b, Karila et al., 2003a). Some of the cardiovascular effects of AAS, such as hypertension, dyslipidemia, and coagulation abnormalities, remit after AAS use was discontinued, but effects such as atherosclerosis and cardiomyopathy appear to be irreversible (Hartgens and Kuipers, 2004, Sullivan et al., 1998b).

Bodybuilders, examined a mean of several years after last AAS exposure, still exhibited impaired myocardial function (D'Andrea et al., 2007) which was associated with the duration and dose of previous AAS use. These results were confirmed in two small cohorts (Krieg et al., 2007, Nottin et al., 2006) where AAS had significantly impaired myocardial function. Two cases of sudden cardiac death were reported in healthy bodybuilders who chronically used AAS (Fineschi et al., 2007), and several case reports document myocardial infarction and stroke in AAS abusers which were partial (Frankle et al., 1988a, Kennedy and Lawrence, 1993, McNutt et al., 1988). In a 12 years follow-up study (Parssinen et al., 2000), the mortality in 62 Finnish power lifters, strongly suspected of having used mega doses of AAS over several years, was 12.9% (mean age at death 43 years) compared with 3.1% in the control group of 1094 subjects (mean age not documented). Suicide and acute myocardial infarction accounted for six out of eight deaths.

Despite the growing number of anecdotal reports of death attributable to apparent cardiac problems among young AAS users (Kanayama et al., 2008), there is no epidemiological evidence for cardiovascular disease due to AAS use (Parssinen and Seppala, 2002, Santora et al., 2006), so that this causality remains to be established. Note, however, that the risk of cardiovascular complications may also be due to the use of other doping drugs, like growth hormone or EPO (erythropoietin).

7.3.4 *Other diseases*

AAS-induced hypogonadism can lead to impaired sexual functioning (Brower, 2002, Pope and Brower, 2009) and infertility (De la Torre et al., 2004c, Menon, 2003b, Turek et al., 1995), but the function normally recovers upon cessation of use.

8 Benzodiazepines

8.1 Acute adverse effects

Benzodiazepines have a dose-dependent respiratory depressant effect, modestly reduce blood pressure and increase heart rate as a result of decrease of systemic vascular resistance (Oikkola and Ahonen, 2008). Following abuse of benzodiazepines, virtually no side effects are seen (Schuckit, 2000) because of the fairly wide therapeutic index. Only occasionally benzodiazepines have been associated with lethal overdoses when used alone (Drummer and Ranson, 1996), but especially when combined with other CNS depressants, such as alcohol (Koski et al., 2002, Serfaty and Masterton, 1993) and opioids including buprenorphine (Tracqui et al., 1998, Kintz, 2001). Very high doses may be fatal due to respiratory depression.

8.2 Chronic adverse effects

No physical effects of benzodiazepines have been reported. Benzodiazepines, especially in combination with other drugs, impair car driving skills, especially in the first days to weeks of treatment. Chronic use will lead to tolerance to many of these impairing effects unless high doses are used (Leung, 2011).

8.3 Disease

Benzodiazepine use has been associated with sexual dysfunction in both sexes, manifested as decreased sexual desire, erectile dysfunction, inhibited orgasm, and inhibited ejaculation (Uhde et al., 1988, Ghadirian et al., 1992, Fossey and Hamner, 1994). Such effects are, however, scarcely seen and based on prospective data with certain design limitations. These side effects seem to emerge after weeks of use and are likely to subside after dose reduction or cessation of use.

9 Khat

Khat has effects on the central nervous system which resemble those of amphetamine. Due to the bulkiness of the plant, high amounts of the leaves must be consumed to attain blood levels of the active ingredient cathinone, which may become harmful. If used in very high quantities, khat (intoxication) may result in cardiovascular toxicity with hypertension and tachycardia, but severe hypertension has not been observed (Hassan et al., 2000, Luqman and Danowski, 1976b).

9.1 Acute adverse effects

The main acute toxic effects of khat use include (cf. Table 1) increased blood pressure, tachycardia, insomnia, anorexia, constipation, general malaise, irritability, migraine and impaired sexual potency in men (Nencini and Ahmed, 1989).

Table 1. Acute effects, but not toxic effects per se, due to khat use

relief of fatigue, increased alertness, reduced sleepiness
mild euphoria and excitement; improved ability to communicate, loquacity
tachycardia, hypertension
moderate hyperthermia
mydriasis, blurred vision
anorexia, dry mouth
constipation (amphetamine-like effect, but supposedly also due to tannins)
psychotic reactions at high doses
irritability and depressive reactions at the end of a khat session
lethargy and sleepy state (next morning)

9.2 Chronic adverse effects

9.2.1 Overview

Khat use is associated with a variety of adverse effects (cf. Table 2).

Table 2. Chronic adverse effects of khat use

Mild depressive reactions during khat withdrawal or at the end of a khat session
Psychotic reactions (hear voices, paranoid state) following frequent use of high doses
Malnutrition
Irritative disorders of the upper gastro-intestinal tract (gastritis, enteritis)
Cardiovascular disorders
Haemorrhoids
Impaired male sexual function, spermatorrhoea, impotence, lower libido
Periodontal disease, mucosal lesions (keratosis)
Genotoxicity, reproduction toxicity, carcinogenicity

9.2.2 Cardiovascular complications

Khat chewing may be a precipitating factor for myocardial infarction as a result of catecholamine release. As compared to non-chewers, khat chewers presenting with acute myocardial infarction were more likely to be young and without cardiovascular risk factors, and were more likely to present during or immediately after khat-chewing sessions (Al Motarreb et al., 2002b).

During khat sessions the incidence of acute myocardial infarction presenting between 2:00 pm and midnight was increased (Al Motarreb et al., 2002a) and khat chewing was reported to have a 39-fold increased risk for acute myocardial infarction (Al-Motarreb et al., 2005). Khat chewing is also a significant risk factor for acute cerebral infarction (Mujili et al., 2005). The prevalence of high blood pressure was significantly higher in khat chewers. Another cardiovascular complication of khat chewing is the higher incidence of haemorrhoids and haemorrhoidectomy found in chronic khat chewers (62% and 45%) as compared to non-khat users (4% and 0.5%) (Al Hadrani, 2000).

A recent study (Ali et al., 2010) in 8176 patients, mainly of Yemeni origin (11% were khat chewers) showed that after adjustment of baseline variables, khat chewing was an independent risk factor for stroke (OR 2.7; 1.3 - 5.9; P=0.01). Cigarette smoking was more prevalent in khat chewers.

9.2.3 *Oral and gastro-intestinal complications*

As a consequence of its mode of consumption khat affects the oral cavity and the digestive tract. A high frequency of periodontal disease has been suggested as well as gastritis (Kennedy et al., 1983), and chronic recurrent subluxation and dislocation of the temporomandibular joint (Kummoona, 2001).

Epidemiological studies, however, have yielded conflicting results. Several studies indicated no such detrimental effects of khat chewing and suggested beneficial effects on the periodontium (Hill and Gibson, 1987a, Jorgensen and Kaimenyi, 1990). Another study could not show a significant role of khat chewing and suggested bad oral hygiene as a major factor in periodontal disease (Mengel et al., 1996). No significant association could be found between khat chewing and oral leukoplakia (Macigo et al., 1995). Khat chewing seems to improve gingival health (Al-Hebshi and Skaug, 2005). Oral keratotic lesions at the site of chewing (Ali et al., 2004b) and plasma cell gingivitis (allergic reaction to khat) (Marker and Krogdahl, 2002) have been reported. The tannins present in khat leaves are held responsible for the observed gastritis (Halbach, 1972, Pantelis et al., 1989). Associations between khat use versus gastric ulcers and constipation has been observed, but its causality is not known (Luqman and Danowski, 1976a).

9.2.4 *Growth retardation*

Retardation of growth rate was considered to be due to decreased absorption of food and not due to decreased food consumption. In pregnant rats, khat reduces food consumption and maternal weight gain, and also lowers the food efficiency index (Islam et al., 1994). Khat extracts and (-)-cathinone produce anorectic effects in animals (WHO, 1980) which is qualitatively similar to that evoked by amphetamine (Goudie, 1985, Zelger et al., 1980).

9.3 **Diseases**

9.3.1 *Cancer disease*

With the micronucleus test to determine genetic damage, an eightfold increase in micronucleated buccal mucosa cells (but not bladder mucosa cells) was seen among khat chewing individuals (Kassie et al., 2001a). Here, khat, tobacco and alcohol showed additive effects, suggesting that khat consumption, especially when accompanied by alcohol and tobacco consumption, potentially causes oral malignancy (Kassie et al., 2001b).

Makki (Makki, 1975) stressed the importance of khat when she found that most of the oral squamous cell carcinomas of her study patients were located in the buccal mucosa and lateral sides of the tongue, which comes into direct contact

with the khat during chewing. Of the 28 head and neck cancer patients in Saudi Arabia (Soufi et al., 1991), ten patients had a history of khat chewing. All were non-smoking chewers and all of them had used khat over a period of 25 years or longer; eight of these ten presented with oral cancers. In some cases the malignant lesion occurred at exactly the same site where the khat bolus was held. The authors concluded that a strong correlation between khat chewing and oral cancer existed. In another study performed in Yemen, 30 of 36 patients suffering from squamous cell carcinoma (in the oral cavity: 17; oropharynx: 1; nasopharynx: 15; larynx: 3) were habitual khat chewers from childhood (Nasr and Khatri, 2000).

9.3.2 *Periodontal disease*

Half of khat chewers develop oral mucosal keratosis of the oral buccal mucosa (Hill and Gibson, 1987b) which is considered as a pre-cancerous lesion that may develop into oral cancer (Goldenberg et al., 2004). Ali et al. reported that 22.4% of khat chewers had oral keratotic white lesions at the site of khat chewing, while only 0.6% of non-chewers had white lesions in the oral cavity (Ali et al., 2004a). The prevalence of these lesions and its severity increased with frequency and duration of khat use. In human leukaemia cell lines and in human peripheral blood leucocytes, khat extract, cathinone and cathine produced a rapid and synchronized cell death with all the morphological and biochemical features of apoptotic cell death (Dimba et al., 2004).

9.3.3 *Reproductive effects*

In man, khat use during pregnancy is associated with lower birth weight. No teratogenic effects have been reported, but detailed studies on the effects of khat on human reproduction are lacking. However, the available data suggest that chronic use may cause spermatorrhoe and may lead to decreased sexual functioning and impotence (Halbach, 1972, Mwenda et al., 2003). In chronic chewers, sperm count, sperm volume and sperm motility were decreased (el Shoura et al., 1995, Hakim, 2002). Deformed spermatozoa (65% of total) have been found in Yemenite daily khat users, with different patterns including head and flagella malformations in complete spermatozoa, aflagellate heads, headless flagella, and multiple heads and flagella (el-Shoura et al., 1995). In rodents, orally administered khat extract induced dominant lethal mutations (Tariq et al., 1990), chromosomal aberrations in sperm cells (Qureshi et al., 1988b), and teratogenic effects (Islam et al., 1994).

10 LSD

On weight basis, LSD is 100 times more potent than the 'magic mushroom' constituents psilocybin and psilocin.

10.1 Acute adverse effects

The earliest effects, seen an hour or so after consuming LSD, are likely to involve stimulant-like physical changes such as pupillary dilation, and increases in heart rate, blood pressure, and body temperature (Abraham, 2004, Pechnick and Ungerleider, 2004). At this stage tremors and paresthesias are likely to occur, along with an increase in blood sugar and hormones, like cortisol, ACTH, and prolactin (Pechnick and Ungerleider, 2004, Strassman et al., 1996). Physical effects of LSD may further include goose bumps, euphoria, uterine cramps and contractions, numbness, muscle weakness, trembling, jaw clenching, impaired motor skills and coordination, nausea, perspiration, saliva and mucous production, sleeplessness and tremors and, occasionally, seizures. These short-term effects appear soon after a single dose and disappear within a few hours or days.

Of drug-related visits to emergency departments in the USA, only 0.1% involved LSD, whereas cocaine was involved in 20%. Deaths due to LSD overdose are seldom reported. Patients generally show sympathomimetic side effects, like mydriasis, hypertension, flushing, tachycardia, and hyperthermia (rarely). Behaviour can be agitated or withdrawn. Adverse reactions are usually seen in inexperienced users or in people who have taken the drug unknowingly. An unexpected stressful setting can cause an acute panic reaction, even in experienced users.

10.2 Chronic adverse effects

Examining nearly a hundred papers, LSD was found to be a weak mutagen, effective only at very high doses. LSD is not carcinogenic and did not cause chromosome damage in human beings at normal doses. The few available prospective studies, mostly of psychiatric patients before and after LSD use, showed no chromosome damage. There was no evidence of a high rate of birth defects in children of LSD users (Dishotsky et al., 1971). This paper is well known and adequately covers the research up to 1971; later studies have allayed persisting doubts.

10.3 Diseases

Those who are tripping on LSD are prone to personal injuries including suicides and fatal accidents.

Although LSD is considered relatively safe when compared with other drugs of abuse, there are case reports of hyperthermia, respiratory failure, and coagulopathies following massive doses (Klock et al., 1974a). Long-time LSD use was suspected to result in chromosomal damage, but this has been consistently refuted; i.e. LSD is not teratogenic (Cohen and Shiloh, 1977). LSD, however, does induce uterine contractions which could disrupt pregnancy. The main reasons of medical support to LSD users are "the bad trip", 'flashbacks', and persistent psychosis. No somatic disease has been reported.

11 Magic mushrooms

11.1 Acute adverse effects

In general, the physiological side effects are not significant and may include dizziness, nausea, weakness, muscle aching, shivering, abdominal pain and dilation of pupils (mydriasis). A UK clubbing magazine survey conducted in 2005 found that over 25 per cent of those who had used hallucinogenic mushrooms in the last year had experienced nausea or vomiting (Mixmag, 2005). Tachycardia is a common finding in patients intoxicated by *Psilocybe* mushrooms. Mild-to-moderate increase in breathing frequency, heart rate (tachycardia of 10 b.p.m.) and systolic and diastolic blood pressure increase (+25, and +10 mmHg, respectively) is observed at 0.2 mg/kg psilocybin p.o. (Gouzoulis-Mayfrank et al., 1999), confirming previous data following an intake of 8-12 mg psilocybin p.o. (Quetin, 1960). Generally, body temperature remains normal, but pronounced physical symptoms such as severe stomach pain, persistent vomiting, diarrhoea etc. have been recorded. The tendency for a temporarily increased blood pressure may also be a risk factor for users with cardiovascular conditions, especially untreated hypertension (Hasler et al., 2004). Acute toxicity of psilocybin is believed to be low, so fatal intoxications related to consumptions of hallucinogenic mushrooms are rare.

11.2 Chronic adverse effects

Though systematic research has not been performed, there is no evidence of chronic toxicity so far. Depending on the setting, the psychological well-being of the user and the dose, intoxications (bad trips) may occur. Very serious intoxications, like severe paranoia, flash-backs, psychosis-like states may lead to accidents, self-injury or suicide attempts. Such severe accidents, however, seldom occur.

11.3 Diseases

No somatic disease related to the use of magic mushrooms has been reported.

12 Ketamine

Ketamine is an anaesthetic with a good safety profile; the main effects being neurobehavioral in nature. The major drawback, which limits clinical use, is the occurrence of emergence reactions in patients awakening from ketamine anaesthesia. Moreover, ketamine differs from other anaesthetics in that it is a cardiovascular stimulant: it increases heart rate, cardiac output and blood pressure. These effects pose no problem except when (dosed to patients or) taken by users with significant ischemic heart disease, high blood pressure or cerebrovascular disorders.

12.1 Acute adverse effects

Low doses of ketamine causes stimulant effects with a temporary increase in blood pressure and heart rate, as well as diplopia and nystagmus (Harari and Netzer, 1994). Tachycardia and hypertension are the most common physical findings after illicit use (Weiner et al., 2000a).

Overdose is rare, but adverse physical effects include hypertension, tachycardia, chest pain and, in more severe reactions, respiratory collapse or heart failure. Less frequently mentioned adverse effects are myalgia, rhabdomyolysis, raised body temperature, hepatic crisis, and mydriasis (Arditti, 2000, Dalgarno and Shewan, 1996, Reich and Silway, 1989a, Siegel, 1978, Weiner et al., 2000b). Rhabdomyolysis may result from muscle rigidity combined with exertion in severe agitation. Large doses result in deep anesthesia with coma and respiratory depression (Reich and Silway, 1989b).

Ketamine induced respiratory depression and cardiovascular pathology are usually rarely serious when only ketamine is used, but may become more serious when ketamine is used in combination with respiratory and central nervous system depressants like ethanol, opioids, barbiturates and benzodiazepines and cardiovascular stimulants, such as amphetamine, ephedrine and cocaine (Buck and Blumer, 1991, Kopman, 1972, Moore et al., 1997)

12.2 Chronic adverse effects

There are no studies which specifically address chronic toxicity of recreational use of ketamine and concomitant abuse of other drugs. Ketamine has not been associated with genotoxicity, carcinogenicity or reproductive toxicity.

12.3 Diseases

Following long-term recreational use of ketamine gastro-intestinal toxicity, particularly abdominal pain (K-cramps) (Jansen, 2000) with unknown aetiology (Muetzelfeldt et al., 2008c) and abnormal liver function (Poon et al., 2010) was reported. Urologic insults due to regular ketamine use were recently reviewed by Smith (Smith, 2010) and Kalsi et al. (Kalsi et al., 2011). Urinary tract symptoms in ketamine abusers are increasingly recognized; 20–30% of frequent users report bladder symptoms (Muetzelfeldt et al., 2008b, Chu et al., 2008c). Case series demonstrate a temporal link between ketamine use (abuse) and urological symptoms, urinary tract damage and renal impairment (Shahani et al., 2007b, Muetzelfeldt et al., 2008a, Chu et al., 2008b), with some but not all improving on cessation of ketamine.

The observation that ketamine may lead to significant urological side-effects was first recognized in chronic ketamine recreational users (Oxley et al., 2009,

Colebunders and Van Erps, 2008, Chu et al., 2008a). Cottrell and collaborators reported increasing numbers of patients chronically using ketamine with urological complications (Cottrell and Gillatt, 2008). Urological side effects to ketamine have been reported in the last year as an emerging problem amongst the drug user population (Shahani et al., 2007a, Chu et al., 2008d).

13 GHB (Gamma Hydroxy Butyric acid)

Bodybuilders have used GHB to increase muscle mass. In a small study conducted in six male human volunteers, GHB (2.5 g) significantly increased prolactin and growth hormone secretion (Takahara et al., 1997). GHB is also used as a sexual adjunct to enhance libido and sexual function, by both heterosexuals and homosexuals (Sumnall et al., 2008).

13.1 Acute adverse effects

Following a typical 65 mg/kg intravenous dose of GHB, sleepiness can occur within five minutes, followed by a comatose state lasting for one to two hours or more, after which there is a sudden awakening. The same dose can also cause hypotonia, bradycardia, nausea, vomiting, random clonic movements of the face and extremities and Cheyne-Stokes respiration (Laborit, 1964, Chin et al., 1992). Physical signs of GHB are bradycardia, respiratory depression and apnoea (Kam and Yoong, 1998, Chin et al., 1998).

13.2 Chronic adverse effects

In rats receiving 170 mg/kg GHB (35 mg per rat) daily for 70 days, no toxicity at organ level (weight, bone marrow, liver and kidneys) was observed during chronic toxicity tests. Symptoms seen in animals treated with high doses of GHB (>35 mg) include various degrees of sleep, bradycardia, decreased body temperature, seizures/spasms, and ultimately respiratory depression, which is fatal (Laborit, 1964). Over half of all patients who present with GHB intoxication (coma) have abused other drugs, as well (Van Sassenbroek et al., 2003, Sporer et al., 2003). The combined use of GHB and alcohol was frequently mentioned. No animal or human data are available concerning reproductive toxicity, neurotoxicity, mutagenicity and the carcinogenic potential of GHB.

13.3 Diseases

No somatic disease has been described for GHB use. The long-term effects are, however, not well known.

14 Ecstasy

14.1 Acute adverse effects

Most often reported effects of ecstasy are (in order of frequency) lack of appetite, jaw clenching, dry mouth, thirst, restlessness, palpitations, impaired balance, difficulty in concentration, dizziness, feeling and sensitivity to cold, drowsiness, nystagmus, hot flashes, trismus, muscular tension, weakness, insomnia, confusion, anxiety, and tremor.

MDMA can also produce panic attacks, delirium, and brief psychotic episodes that usually resolve rapidly when the drug action wears off. Short-term side effects (up to 24 hours after ecstasy consumption) most often reported are (in order of frequency) fatigue, heavy legs, dry mouth, loss of appetite, insomnia, drowsiness, weakness, muscular tension, lack of energy, difficulty concentrating, and headache. Late short-term residual side effects (up to seven days after ecstasy use) include fatigue, irritability, anxiety, lack of energy, depressed mood, insomnia, drowsiness, and muscular tension (De la Torre et al., 2004b). Severe intoxication can include delirium, coma, seizures, hypotension, arrhythmias, hyperthermia (>40°C), and renal failure associated with rhabdomyolysis. A serotonin syndrome (increased muscle rigidity, hyperreflexia, and hyperthermia) and intracranial haemorrhage have been described. Hyperthermia may result from a direct action of the drug on the CNS temperature regulating centre and vasoconstriction of skin vessels and can be related to muscular activity associated with dance or tremor and rigidity, high ambient temperatures in crowded places, and dehydration. Heat stroke is a severe complication that can cause death; it includes hyperthermia, rhabdomyolysis, myoglobinuria, disseminated intravascular coagulation, and renal failure. Hyponatraemia is an uncommon complication associated with excessive water intake; the syndrome of inappropriate antidiuretic hormone (SIADH) is usually present, with increased levels of anti-diuretic hormone (ADH).

14.2 Chronic adverse effects

Fulminant hepatitis and hepatic necrosis have been described (De la Torre et al., 2004a). Furthermore, no major chronic effects have been reported (except from those resulting from overdosing), mild negative effects on concentration, cognition, and sleep have been reported.

14.3 Diseases

MDMA use is not associated with frequently occurring serious disease. Long dance marathons are often associated with MDMA use. Specifically when used in the setting of crowding and vigorous dancing, such as in 'raves' or clubs, MDMA may lead to volume depletion, body temperatures greater than 40°C, cardiovascular collapse, and convulsions. Other symptoms are rhabdomyolysis, disseminated intravascular coagulation, and acute renal failure (Gouzoulis-Mayfrank and Daumann, 2006, Mccann et al., 1996, Williams et al., 1998) (Pechnick and Ungerleider, 2004). This is thought to be due to the combination of sympathomimetic effects including cutaneous vasoconstriction and extreme physical exertion in hot and poorly ventilated conditions, although some features are those of serotonin syndrome (Mueller and Korey, 1998). Physical effects include impairment of balance, conjunctival infection, increased heart rate, orthostatic hypotension, peripheral vasoconstriction with cold extremities, dry

mouth, and increased appetite (Ashton, 2001c, Hall and Solowij, 1998, Hall and Degenhardt, 2009).

Such manifestations are related to the life-threatening serotonin syndrome, which is characterized by muscle rigidity, shivering, tremor and increased deep tendon reflexes. The excessive muscle contraction leads to hyperthermia with an associated mortality rate of 10 to 15% (Hall and Henry, 2006). In 1999, the risk of death for first-time users of ecstasy was estimated to be between 1 in 2,000 and 1 in 50,000 (Gore, 1999). These data are consistent with Dutch data, reporting one to three fatal cases per annum. However, considering the high prevalence of use of ecstasy, fatal incidents following ecstasy occur rarely. MDMA ingestion may lead to hepatotoxicity, including hepatic failure requiring transplantation, (Brauer et al., 1997, Garbino et al., 2001, Henry, 1992, Jones and Simpson, 1999, Milroy et al., 1996, Sano et al., 2009). MDMA induced liver failure is likely to be mediated by a hypersensitivity reaction (Andreu et al., 1998, Ellis et al., 1996, Fidler et al., 1996). Moreover, liver hypofunction due to the use of certain pharmaceutical drugs or hepatitis is an absolute contra-indication of ecstasy use.

15 Cannabis

15.1 Acute adverse effects

Dose-dependent, THC increases in heart rate, which via a feedback compensation mechanism leads to hypotension. Due to the decrease in vascular resistance and the increase in blood flow to the limbs, orthostatic hypotension develops, which leads to dizziness (in relatively inexperienced users). Carbon monoxide in the smoke may increase dizziness. Bradycardia develops due to tolerance. Secondary effects of THC, other than those on the central nervous system and circulation are: hyporeflexia, miosis or mydriasis and hyperemia of the conjunctiva, bronchodilation, dry mouth, nausea, vomiting, diarrhoea, abdominal cramps, tremors, muscle weakness and urinary retention. Severe intoxications lead to depression of the central nervous system and sometimes coma.

15.2 Chronic adverse effects

15.2.1 Pulmonary system

Cannabis smoke condensate ('tar') is positive in the Ames test, scoring mutagenicity (Busch et al., 1979, Sparacino et al., 1990). The carcinogenic potential of cannabis smoke is weak as compared to tobacco smoke condensate (Hoffmann et al., 1975), whereas cannabinoids themselves are not mutagenic (Hall et al., 2005) or carcinogenic (Chan et al., 1996). Some cannabinoids are even anti-carcinogenic (Guzman, 2003), but their clinical significance is unclear. Ashton (Ashton, 2001d), however, showed that marijuana reefers contain 3–5 times more tar and carcinogens as compared to cigarettes. In the tissues and cells of the airways of cannabis smokers, the same pathological changes are found as in tobacco smokers, matching early stages of bronchial carcinoma (Barsky et al., 1998).

15.2.2 Immune System

The impairment of immune function by THC may increase the risk for infections and cancers (Ibrahim et al., 2003, Quartilho et al., 2003, Vigano et al., 2003). Excluding the effects in the lungs, there is, however, little evidence for a reduced resistance to infections in cannabis users (Kraft and Kress, 2004). Reduced pulmonary resistance may also result from smoking mouldy cannabis (Sakkour et al., 2008).

15.2.3 Endocrine System

Decreased levels of hormones, including luteinizing hormone and testosterone, have been demonstrated in heavy marijuana smokers (Block et al., 1991), but these abnormalities appear to be temporary. It is also possible that growth hormone production and prolactin levels may be altered in heavy marijuana smokers, but the clinical significance to humans has not been established (Murphy et al., 1998).

15.3 Disease

15.3.1 Pulmonary disease

Though low marijuana use is hardly harmful, the (medical) harm of marijuana use grows with increasing amount, frequency of intake, and years of exposure to these drugs (Crean et al., 2011, Dewit et al., 2000, Taylor et al., 2002). In

heavy smokers of marijuana a chronic inflammation of the sinuses (nose; sinusitis), as well as pharyngitis (throat), has been reported (Hall, 1998). Large-scale prospective studies document the negative pulmonary effects of marijuana over time (e.g., (Sherrill et al., 1991, Tashkin, 1993a, Tashkin et al., 1997a). Though not fully consistent, the results converge on the observation that greater duration of marijuana use is related to increased bronchitis symptoms (e.g., coughing, wheezing (Tashkin, 1993b, Tashkin et al., 1997b). The pulmonary exposure to carbon monoxide and carcinogens following smoking of cannabis is four to five times higher as compared to tobacco smoking, mainly because cannabis smokers inhale 60% more smoke, inhale the smoke about 30% deeper and hold the smoke four times longer (Wu et al., 1988). An acute effect of THC is bronchial dilation (Tashkin et al., 1973, Tashkin et al., 1974, Tashkin et al., 1975, Calignano et al., 2000). However, exposure to the toxic smoke compounds may result in chronic cough, sputum production, wheezing and acute bronchitis (Taylor et al., 2000, Tashkin et al., 1987, Bloom et al., 1987), which finally gives long-term airway inflammation (Tashkin, 2005). It was estimated that 3 marijuana reefers have the same risk to produce bronchitis and emphysema as 20 cigarettes (Ashton, 2001b). Others (Tetrault et al., 2007) found no consistent relationship between cannabis smoking and lung function, but believe that the risk of COPD is increased by the respiratory symptoms. Interestingly, smoking only marijuana was not associated with an increased risk of COPD (Tan et al., 2009).

15.3.2 *Cancer disease*

Most investigations to the relation between marijuana use and cancer suggest that there is an increased risk of lung cancer among more frequent users of marijuana (Caplan and Brigham, 1990b). Several reviews describe the epidemiological studies on the association between cannabis and lung cancer (Hashibe et al., 2005, Mehra et al., 2006). After correction for confounding variables including cigarette smoking, an increased risk of lung cancer was found in the highest tertile (>10.5 joint-years) (OR=5.7, 95% CI=1.5-21.6) (Aldington et al., 2008). Used as a continuous variable, each joint-year was associated with is low relative risk of 1.08 (CI=1.02-1.15). Two studies in North Africa, where cannabis, in contrast to New Zealand, is smoked mixed with tobacco, gave similar risks (Hsairi et al., 1993, Sasco et al., 2002). Two U.S. studies found no association between cannabis use and cancer (Hashibe et al., 2006, Sidney et al., 1997). Chronic smoking of marijuana appears to be associated with respiratory tract carcinoma and head and neck cancer in young adults (Ho et al., 2009, Tashkin, 2001, Taylor, 1988, Van Hoozen and Cross, 1997, Zhang et al., 1999). Possibly, heavy marijuana smokers have the same increased risk of cancers of the head and neck as heavy tobacco smokers.

It can be concluded that smoking marijuana slightly increases the lung cancer risk, but the risk is smaller than with tobacco smoking. Partly this is due to the generally relatively short period of cannabis use. The risk may be higher for frequent or long-term cannabis smoking. Epidemiological data allow no clear conclusion about other cancers.

15.3.3 *Cardiovascular disease*

Marijuana acutely produces tachycardia (an increase in heart rate), vasodilatation, and a negative inotropic effect (decreased heart contractions) (Ashton, 2001a), which – especially in heart patients – may be harmful due to impaired oxygen supply to the heart muscle. Similarly, the amount of exercise an individual on cannabinoids can tolerate before the onset of heart pain or angina is decreased.

Smoking cannabis increases the risk of myocardial infarction shortly after smoking, and patients with angina pectoris and ischemic cardiovascular conditions (Grotenhermen, 2007, Sidney, 2002) are at higher risk. Ventricular conduction disturbances are occasionally observed which may result in cardiomyopathy (Daccarett et al., 2007). Epidemiological studies indicate no permanent increased risk of cardiovascular disease. Contributing risk factors are male gender, tobacco use and obesity that often exist among cannabis users (Mittleman et al., 2001).

15.3.4 *Reproductive disorders*

Animal studies suggests that heavy marijuana use impairs the reproduction capacity (Hall et al., 1994), but controlled evidence among humans is lacking (Caplan and Brigham, 1990a). Chronic use of cannabis may inhibit the secretion of reproductive hormones and induce impotence in men and menstrual irregularities in women (Hollister, 1986b). Chronic marijuana use may decrease the male prostate and testes, reduce sperm count and block ovulation, These changes are reversible, and do not imply infertility (Abel, 1981).

15.3.5 *Other diseases*

Like tobacco smokers, recreational users of cannabis smokers generally have poorer oral health than non-users, with an increased risk of dental caries and periodontal diseases (Cho et al., 2005). The illicit use of cannabis pose certain obstacles and challenges to the dental professional.(Maloney, 2011).

15.4 **Health benefits**

Although most research has focused on the negative health consequences of marijuana use, the drugs also appears to retain possible health benefits. Marijuana has been shown to decrease intraocular eye pressure, involuntary movement and perceived pain, and to stimulate appetite (Hollister, 1986a, Hollister, 1992a). The strongest evidence of possible health benefits for marijuana use appears to be focused on increasing appetite, and decreasing nausea and vomiting (as a result of cancer therapy), and possibly improving pain tolerance (Hollister, 1992b).

16 Amphetamine

16.1 General remarks on amphetamine and methamphetamine

Amphetamine is a metabolite of methamphetamine, so that methamphetamine users will be exposed to both substances simultaneously. In the Netherlands, amphetamine is the predominant amphetamine and it is usually administered by either the oral or the intranasal route. Intravenous injection is rare.

16.2 Acute adverse effects

Methamphetamine more efficiently penetrates the brain and has a longer half-life than amphetamine; its effects persist for 6 to 24 hours longer than amphetamine. Methamphetamine more potently stimulates CNS with fewer peripheral effects compared to amphetamine (Buchanan and Brown, 1988), though large doses induce hypertension.

Table 3. Routes of administration effects and risks

Route	Effect	Risks
Intravenous	High and rapid peak effect followed by reduction in intensity over the next four to six hours	Needle sharing. Injection risks include: <ul style="list-style-type: none"> • inflammation, infection, scarring, or abscess at i.v. site • acute intoxication risks such as cardiovascular complications (incl. arrhythmias, cerebrovascular accident)
Smoking/ inhalation	Less intense onset and duration of effect	Best route to control the dose, though relatively uncommon. Relatively rapid effect. May result in sore throat, bloody sputum, and exacerbation of asthma.
Snorting	Weaker onset and slower reduction in intensity than i.v. but slightly longer lasting	Damages epithelium and nasal septum, potentially causing nasal ulcers, runny nose, sinusitis, and septum perforation.
Swallowing or 'bombing'	Delayed absorption (about 30 minutes to 'come on', slower peak, slower reduction, lasting around six hrs)	Impatience waiting for effect, inability to control the dose, or seeking a stronger or more intense effect may result in taking more drugs, possibly increasing intoxication risks.
Anal (shelving)	Unpredictable effects, varies with quality and quantity of the drug formulation	Highly acidic forms may irritate mucosal lining. Time is required for absorption to occur before effect is experienced (see oral use above).

From (NCETA, 2004)

Tables 3 and 4 give an overview of differential effects of amphetamine administration through different routes and the potential acute physical effects from using low and high doses of amphetamine (reproduced from an Australian handbook (NCETA, 2004)).

It is evident from Tables 3 and 4 that both route of administration and dose determine the extent of the effects, with a potential for more harmful effects at high doses. Note that the data of Tables 3 and 4 originate from Australia, where methamphetamine is the predominant amphetamine. Compared to alcohol

(2056), cannabis (281) and heroin/cocaine (230), the number of amphetamine-related accidents in Amsterdam is very low (3) (Van Laar et al., 2008).

Table 4. Potential acute physical effects from using low and high doses of amphetamines (NCETA, 2004)

System	Low doses	High doses
Cardiovascular	tachycardia (possibly brief bradycardia), hypertension palpitations, arrhythmias	cardiac stimulation (tachycardia, angina, arrhythmia*, MI) vasoconstriction/hypertension cardiovascular collapse*
Respiratory	increased respiration rate and depth	respiratory difficulty/failure*
Gastrointestinal	nausea and vomiting constipation, diarrhoea or abdominal cramps	dry mouth nausea and vomiting abdominal cramps
Skin	pale sweaty skin, hyperpyrexia	flushing or pallor hyperpyrexia, diaphoresis
Skeletal	increased deep tendon reflexes	-

* *death due to amphetamine overdose; however, death is rare.*

16.3 Chronic adverse effects

16.3.1 Cardiovascular pathology

Kaye and co-workers (Kaye et al., 2007) concluded that there is sufficient clinical and experimental evidence to suggest that methamphetamine has adverse and potentially fatal effects on the cardiovascular system. Yet, it should be noted that the majority of the cases reviewed by Kaye et al. concern amphetamine intoxications, which implies that both amphetamines have the potential to cause cardiovascular harm. Martindale's monograph on dexamphetamine sulphate (Reynolds, 1993) states that cardiomyopathy rarely occurs when amphetamine is used in therapeutic dose, but in acute overdose chest pain, cardiac arrhythmias and circulatory collapse may occur.

16.4 Disease

16.4.1 Cardiovascular disease

The use of stimulants, like amphetamines is associated with cardiac complications (Bashour, 1994, Ragland et al., 1993). Clinically, the effects of amphetamine are very similar to those of cocaine, but the amphetamine effects are more prolonged (half-life of 10 to 15 h). Stroke has been reported in patients with amphetamine intoxication; it results from hypertension and is usually haemorrhagic (Agaba et al., 2002, El-Omar et al., 1996, Petitti et al., 1998). There have also been reports of cerebral vasculitis and haemorrhage with chronic abuse of amphetamine (Buxton and McConachie, 2000, Matick et al., 1983, Shaw et al., 1985). Cardiomyopathy is also seen in chronic amphetamine abuse (Smith et al., 1976). Systemic necrotizing vasculitis, resembling periarteritis nodosa, has been associated with chronic amphetamine abuse (Welling et al., 1998).

In a cohort of discharges from Texas hospitals during 2000-2003, 11,011 cases (aged 18-44 years) of acute myocardial infarctions (AMI) were identified (Westover et al., 2008). Using multiple logistic regression analysis, amphetamine abuse was significantly associated with AMI (odds ratio=1.61; 95% CI=1.24-2.04, p=0.0004; adjusted for cocaine abuse, alcohol abuse, tobacco use, hypertension, diabetes mellitus, lipid disorders, obesity, congenital

defects, and coagulation defects). The population attributable risk suggests that amphetamine abuse is responsible for 0.2% of AMIs in the state of Texas.

16.4.2 Infections

Specific risks associated with intravenous injection are higher HIV seroprevalence due to needle-sharing (Darke et al., 2007a). Poorly maintained injection sites (e.g. infection) may cause callusing, scarring or abscesses; contaminants present (from acute injection or due to longer term accumulation) may result in lung or cardiac emboli, cardiac valve infections, or stroke (NCETA, 2004).

16.4.3 Other diseases

Sexual dysfunction; weight loss, malnutrition, (these may develop to eating disorders, anorexia or nutritional deficiency); lowered immunity, although with re-establishment of self-care and eating habits, likely to resolve over time (NCETA, 2004).

17 Methamphetamine

Related to the effects described above, heavy and long-term chronic methamphetamine use can result in many life-threatening medical illnesses and disabilities (Mooney et al., 2009).

17.1 Acute adverse effects

Methamphetamine use elicits euphoria, and increases in blood pressure, body temperature, heart frequency, and breathing rate. Other acute clinical symptoms include reduction in fatigue, hunger, and an increase in energy, sexual drive, and self-confidence. Depending on the nature and extent of abuse, the physiological effects can include intense bruxism, shaking, disrupted menstrual cycles, stomach cramps, 'formication', or the sensation of insects creeping on the skin, and insomnia (Richards et al., 1999b).

17.2 Disease

17.2.1 Cardiovascular disease

Cardiopulmonary consequences are common among methamphetamine abusers. Methamphetamine use is associated with ischemic stroke, intracerebral and subarachnoid haemorrhage, especially among young patients (Ho et al., 2009). Both ischemic and haemorrhagic stroke has been reported with methamphetamine abuse, and in some cases the stroke was delayed by 10 to 12 hours after last use (Perez et al., 1999, Rothrock et al., 1988). Cardiovascular symptoms, including irregular heart beats, have developed in more than half of the methamphetamine abusers (Beebe and Walley, 1995). Methamphetamine use related cardiomyopathy may be reversible, but this depends on cessation of drug use (Islam et al., 1995). Pulmonary oedema is reported in over 70% of methamphetamine-related deaths (Karch et al., 1999, Kaye et al., 2009).

Methamphetamine overdose related symptoms seen in emergency rooms are chest pain, hypertension, shortness of breath, acute coronary syndrome and tachycardia (Richards et al., 1999a). Acute coronary syndrome occurs in 25% of methamphetamine abusers admitted for chest pain (Turnipseed et al., 2003), possibly resulting from myocardial ischemia and its attendant risk of arrhythmias and cardiogenic shock (Wijetunga et al., 2004). Damage to small blood vessels in the brain can result in stroke, paralysis, and brain damage (Newton et al., 2003, Wang et al., 2004).

17.2.2 Periodontal disease

'Meth mouth' and other oral complications are common among chronic methamphetamine abusers, even though the contribution of level of methamphetamine abuse to their aetiology is questionable. Oral health problems most often seen among methamphetamine abusers include rampant caries, tooth fracture, and periodontal disease (e.g. gingivitis, periodontitis) (Shaner et al., 2006, Shaner, 2002). In addition to caries and gingivitis, methamphetamine abusers often present with tooth wear and temporomandibular joint syndrome related to bruxism, which may be a reaction to anxiety and restlessness, especially during early abstinence (Curtis, 2006, Hamamoto and Rhodus, 2009).

17.2.3 *Dermatological disorders*

Methamphetamine abusers often show skin excoriations, cellulitis, abscesses or cutaneous ulcers, which result from skin scratching following sensations of bugs crawling below the skin (Bostwick and Lineberry, 2006) or injection of methamphetamine. Methamphetamine abusers may show self-inflicted wounds supposed to result from activation of dopaminergic pathways (Israel and Lee, 2002).

The top four medical conditions associated with methamphetamine-related emergency room visits were mental health (18.7%), trauma (18.4%), skin infections (11.1%), and dental diagnoses (9.6%). In the US, the annual costs for methamphetamine-related ED visits (2.4% of all visits) averaged seven million dollar (Hendrickson et al., 2008).

18 Opiates

Adverse side effects from opiates are seen in drug abusers who take an overdose (intentional or unintentional), but also in medical patients who are treated with opiates are depicted in Table 5.

Table 5. Medical somatic complications of opiate use

Respiratory	Cardiovascular	Other
Cough suppression	Hypotension	Pruritis (irritation of the skin) ¹
Hypoventilation	Bradycardia	Progressive nephrotic syndrome ²
Respiratory arrest	Conduction abnormalities (e.g. propoxyphene)	
Pulmonary oedema	Metabolic Hypothermia	
	Cool, moist skin	

- 1. This is very common in patients receiving opiates, as well as in addicts.*
- 2. Chronic abuse of heroin has been associated with progressive nephrotic syndrome resulting in renal failure (Dubrow et al., 1985).*

19 Buprenorphine

19.1 Acute adverse effects

Buprenorphine has the typical side effects of opioids, such as constipation, vomiting, anxiety, sedation, disturbed sleep, drowsiness, dizziness, headache, pruritis, dry mouth, miosis, orthostatic hypotension, sweating, nausea, male ejaculatory difficulty, decreased libido and urinary retention. Respiratory depression may occur following the high doses, but life-threatening respiratory depression is much less likely with partial agonist buprenorphine as compared to heroin and morphine. Hepatic necrosis and hepatitis with jaundice have been reported, especially after intravenous injection of crushed tablets (Boothby and Doering, 2007, Karch, 2006, Vadivelu and Hines, 2007). Studies revealed that a single dose of buprenorphine up to 70 times the recommended analgesic dose did not induce life-threatening effects, indicating a broad therapeutic index.

19.2 Chronic adverse effects

There is no evidence of organ damage with chronic use of buprenorphine, although induction of liver enzymes is sometimes seen. Buprenorphine in the presence of other narcotic analgesics, general anaesthetics, antihistamines, benzodiazepines, phenothiazines, other tranquilizers, sedative/hypnotics or other CNS depressants (including alcohol) may exhibit severe CNS depression.

19.3 Diseases

No specific disease known other than those described for opiates.

20 Methadone

20.1 Acute adverse effects

Opioids, such as methadone, morphine and heroin, show relatively low toxicity when they are used in normal doses. The effects of methadone last longer than heroin and methadone easily accumulates with repeated dosing. It causes respiratory depression probably deeper than heroin. Although it is less sedative, repeated administration of methadone would cause severe sedation in non-tolerant users. At high dose, methadone prolongs the QT interval ('torsade de pointes').

Table 6. Side effects of opioids

Frequency	Symptoms
Common	Nausea, vomiting, constipation, drowsiness, mental confusion
Moderate	sweating, pruritus, dry mouth, hallucinations, dysphoria, urinary retention, headache
Rarely	thrombocytopenia, rash, urticaria, vertigo, palpitations, orthostatic hypotension

Table 7. Side effects of opioids

System	Effects
Cardiovascular	Hypotension, bradycardia, peripheral vasodilation, tachycardia, sinus bradycardia. Two important side effects of methadone are respiratory depression (when overdosed), cardiac arrhythmias and prolongation of the QT interval (Chugh et al., 2008). The membrane stabilizing properties of methadone may cause cardiac arrhythmias or cardiovascular collapse
Liver	Methadone is metabolized by the liver. Ten patients deceased two to six days after the start of a methadone maintenance programme with an average daily dose of 60 mg of methadone (Drummer et al., 1990). Probably the methadone piled in these patients to very toxic levels, because all patients suffered from chronic hepatitis and liver disease
Urogenital	Reduced urinary excretion, and urinary tract spasms of the ureter
Endocrine	Hypoadrenalism in chronic methadone users (Dackis et al., 1982, Pullan et al., 1983) comparable to ACTH deficiency. There are also indications that methadone inhibits the adrenal function.
Gastrointestinal	Nausea, vomiting, constipation, anorexia, stomach cramps, diseases of the bile duct
Neuromuscular	Rhabdomyolysis, weakness overall
Pulmonary	Apnea, respiratory depression, dyspnoea
Miscellaneous	Miosis, histamine secretion, skin rash, hives, itching

20.2 Chronic adverse effects

Methadone affects the sexual function. In male heroin addicts receiving methadone maintenance programme, sexual performance was significantly reduced (Cicero et al., 1975). The function of the secondary sex organs was significantly decreased compared to untreated heroin addicts and non-addicted

controls. Serum testosterone concentrations in the methadone group were some 43% lower.

Clients in a methadone maintenance programme or patients for the treatment of pain regularly administered methadone, primarily suffer from constipation, but also actually have a few other medical problems due to methadone, as long as they continue taking the methadone (Novick et al., 1993). Other complaints are contraction of the pupils (and therefore bad night vision), blurred vision, sweating, decreased libido, menstrual disorders, urinary retention, insomnia, respiratory disorders and occasionally pain in the joints and bones.

Various studies have shown that methadone substitution programmes have a positive impact on the reduction of 'social harm' caused by abuse of opiates, particularly heroin. What is meant here is public health (reducing HIV infections, hepatitis, etc.) (Skeie et al., 2008). Also, the health of the individual patient is generally improving (Ball et al., 1988, Caplehorn et al., 1996b, Ward et al., 1992, Ward et al., 1999) and mortality among opiate users is decreased (Caplehorn et al., 1996a, Risser et al., 2001). Methadone maintenance programmes appear to be cost effective (Connock et al., 2007), because they reduce health care costs.

21 Heroin

Illegal use of heroin is usually accompanied by a number of adverse health effects, which often result from the poor social and hygienic conditions in which many users find themselves. Dirty syringes from one to the other user are passed and the impurities in street heroin can be purchased for serious health problems.

In the Netherlands, most heroin addicts prefer to smoke the drug by 'chasing the dragon', but in many other countries syringes are still the most common route of administration. It is assumed that 'chasing the dragon' is a safer method than injecting heroin. The population of heroin users in the Netherlands is aging, which is associated with an increase in health problems in this population (NDM, 2011). How many of the heroin users are problem users is hard to determine. It is clear that problem opiate users are all using cocaine as well. In low to moderate doses, the adverse effects of heroin are relatively mild (e.g., nausea and vomiting commonly occur). Gradually, the user develops tolerance for side effects and the desired effects are more appreciated. Tolerance, however, usually occurs quickly for the desired effects such as euphoria, analgesia and rest. As compensation for the development of tolerance to increase, users usually take their daily dose, and/or change to another mode of administration, so that more of the substance reaches the brains and the effect occurs more rapidly.

21.1 Effects of short-term use

21.1.1 Use at lower doses

Table 8. Effects of short-term heroin use in low dose

Respiratory	Slowing of the respiratory rate, which becomes more pronounced at higher doses.
Gastrointestinal	Nausea and vomiting (very common among inexperienced users), reduced appetite, decreased gastric motility and constipation
Other	Reduced libido, itching or burning sensation on the skin, increased urinary output, slightly lowered body temperature, sweating

21.1.2 Use at higher doses

With administration of higher doses intensification of the low-dose effects of heroin occurs and lasts longer. At higher dose, the sensitivity and emotional response to painful stimuli decrease, the ability to concentrate is impaired, the probability of sleep increases, breathing becomes progressively slower and more shallow, heart rate gradually slows and blood pressure decreases.

The most desired effect, referred to as a 'rush', occurs almost immediately following inhalation or intravenous injection, and the subjective experience has been described as akin to an intense orgasmic sensation in the abdomen.

21.2 Chronic adverse effects

Pulmonary oedema after overdose is a common cause of death among heroin addicts. Nausea and hypotension seem to occur less frequently than with the use of morphine (Haemmig and Tschacher, 2001). Heroin inhalation impairs pulmonary function and may lead to pulmonary oedema (non-cardiogenic). Heroin depresses the respiratory centre, releases histamine (which can trigger

asthma), and results in septic emboli and pulmonary oedema, or 'heroin lung'. Pulmonary oedema is a serious complication, which may be due to release of histamine, with increased pulmonary lymph flow and capillary permeability (Dettmeyer et al., 2000). Bilateral pulmonary oedema associated with heroin abuse has been reported several times (Anderson, 1986). Bronchospasm has been noted following the use of street heroin, perhaps due to contaminants (Schoer and Groden, 1999).

Adverse physical health effects directly related to long-term heroin use are reduced libido, constipation, pupillary constriction (which adversely affects night vision), menstrual irregularity and certain types of respiratory impairment.

21.3 Disease following chronic heroin use

The disease burden of heroin use is significant and is associated with premature death from drug overdoses, violence, suicide, alcohol-related causes, HIV-infection, liver disease and renal disease (Hall et al., 2006). Heroin-associated physical disease may result from the use of adulterated heroin, the use of non-sterile injection needles, or the habit of smoking heroin. In addition, heroin dependence is associated with unsafe sex practices with the concomitant risk of acquiring a sexually transmittable disease.

In 2009 in the Netherlands, the primary diagnosis in hospital-admitted opiate users (abuse or dependence as the second diagnosis) was respiratory disease (29%), accident injury (13%), alimentary tract disease (8%), intoxication (5%), abuse or dependence of alcohol and/or other drugs (9%), and skin disease (3%) (NDM, 2011).

21.3.1 Pulmonary disease

Although inhaling is considered to be a safer mode than injecting heroin, it is not without risks. Overdose mortality may occur (Darke and Ross, 2000) and in individual cases, lethal leucoencephalopathy has been reported (Kriegstein et al., 1997, Wolters et al., 1982). Contamination of small batches of drug by an unknown substance may be responsible for this, because the observed cases are locally clustered. Heroin affects the respiratory control centres which may lead to fatal pulmonary depression (White and Irvine, 1999). Like other opiates, heroin is known to release histamine (Edston and Hage-Hamsten, 1997, Withington et al., 1993a), and asthma triggered by inhalation of heroin has been reported (Cygan et al., 2000, Hughes and Calverley, 1988, Withington et al., 1993c). Buster et al., 2002 (Buster et al., 2002, Withington et al., 1993b) found that chronic heroin use by chasing the dragon is related to impaired lung function and higher prevalence of dyspnoea, although they could not exclude that part of the observed lung function impairment was caused by tobacco smoking.

21.3.2 Infections

Illicit opiate injectors show highly elevated prevalence levels of blood-borne infectious disease, specifically HIV and hepatitis C (Fischer et al., 2006). Prevalences may vary considerably between populations. For example, HIV prevalence levels in Canadian injecting drug populations ranged from 10% to 35%, whereas hepatitis C infection levels ranged from 40% to 90% (Fischer et al., 2004, Zou et al., 2000). In the Netherlands, HIV incidence associated with injecting drug use is relatively low. In 2009, less than 1% of newly diagnosed HIV infections were associated with injecting drug use (NDM, 2011). Infections at injection sites and collapsed veins resulting from repeated injections are very common. Tetanus, viral hepatitis (which is a common cause of regular heroin users' admission to hospital medical services), acquired

immune deficiency syndrome (AIDS), endocarditis and other pulmonary complications including tuberculosis all occur more frequently among heroin addicts than in the general population.

21.3.3 *Renal disease*

As reviewed by Jaffe (Jaffe, 1983), previous studies identified a spectrum of renal diseases in heroin users. Some patients develop renal insufficiency after intravenous heroin use (Rice et al., 2000). Its aetiology is unclear. The predominant renal lesion in black heroin users is focal segmental glomerulosclerosis and in white heroin users membranoproliferative glomerulonephritis that may result in renal failure (Cunningham et al., 1983). It is not known whether heroin is causal in the development of renal disease or whether the disease is related to socioeconomic, genetic, cultural or behavioural factors. No well-designed prospective epidemiological studies exist that have assessed the incidence and prevalence of renal disease in heroin users (Jaffe and Kimmel, 2006b).

21.3.4 *Other diseases*

Although there is no evidence of permanent CNS impairment specifically due to chronic heroin use, diminished oxygen supply resulting from a large overdose may result in brain damage. Malnutrition, poor housing, untreated illness and frequent use of and physical dependence on other psychoactive drugs, together with heroin dependence, are likely to result in a generally poor condition and lowered resistance to disease.

21.4 **Mortality**

In Europe, between 10% and 23% of mortality among those aged 15 to 49 might be attributed to opiate use (Bargagli et al., 2006, Bloor et al., 2008, EMCDDA, 2010). Heroin users have a death rate 13-17 times that of their age-matched peers, and a 14-fold risk of suicide (Darke et al., 2007b, Darke et al., 2010, Hickman et al., 2003, Lind et al., 1999, Sporer, 1999). Between 10% and 35% of deaths in heroin-dependent individuals are due to suicide. In the Netherlands, death by opiate overdose is relatively low (52 cases in 2008) (NDM, 2011). Death among opiate-dependent users is mostly caused by lung disease, liver disease (cirrhosis), infectious disease (endocarditis, HIV) and violent acts.

Overall mortality rate for any liver disease in an heroin-dependent Australian ageing cohort was 1.4 deaths per 1,000 person-years (95% CI: 1.1-1.7), 17 times higher than in the general population (95% CI: 13.4-21.3). Liver mortality increased over time and became the most common cause of death in this ageing heroin-dependent cohort (Gibson et al., 2011).

21.5 **Economical burden**

In the United States, the costs of medical care for heroin addiction has been estimated to be approximately € 3.6 billion in 1996 (Mark et al., 2001). Globally, illicit opiate use was estimated to account for 0.7 per cent of global disability-adjusted life years (DALYs) in 2000 when estimates of morbidity attributable to illicit drug use were added in (WHO, 2003). This percentage is probably an underestimate of the disease burden attributable to illicit opioids, because they omit differences across subregions in the quality of data on causes of mortality and estimates of mortality and morbidity attributable to hepatitis and violence (Degenhardt et al., 2004).

22 Cocaine

22.1 Acute adverse effects

Cocaine is a sympathetic stimulant, because it inhibits the neuronal reuptake of catecholamines. Acute cocaine administration increases heart rate, vasoconstriction and hypertension, and hyperthermia. At low doses, these physiological changes are usually not harmful, but they can be toxic or even fatal at high dose. Adverse consequences of heavy cocaine use are seizures, heart failure, stroke, or intracranial haemorrhage (Brands et al., 1988). Cocaine is vasoconstrictive and has local anaesthetic effects and type I antidysrhythmic properties (Bauman et al., 1994, Winecoff et al., 1994). The initial effect of cocaine on the cardiovascular system is a transient bradycardia, secondary to stimulation of the vagal nuclei. Tachycardia results from increased central sympathetic stimulation. The combined use of cocaine and alcohol generates the metabolite cocaethylene, which has a direct myocardial depressant effect (Henning et al., 1994) that is independent of coronary artery vasoconstriction (Pirwitz et al., 1995). The combination of cocaine and tobacco smoking gives coronary vasoconstriction in a synergistic manner (Moliterno et al., 1994).

The severe effects depicted in Table 9 are mostly seen with high-dose use, particularly in individuals with long-standing patterns of chronic intake. Some of these aversive effects (e.g., irritability) are present in most high-dose users, whereas others mainly occur in cases of cocaine-induced psychosis (e.g. incoherence or delusions; see the paragraph on psychiatric effects).

Table 9. Behavioural and subjective effects of cocaine in humans (Meyer and Quenzer, 2005)

Mild to moderate effects	Severe effects
Mood amplification; both euphoria and dysphoria	Irritability, hostility, anxiety, fear, withdrawal
Heightened energy	Extreme energy or exhaustion
Sleep disturbance, insomnia	Total insomnia
Motor excitement, restlessness	Compulsive motor stereotypies
Talkativeness, pressure of speech	Rambling, incoherent speech
Hyperactive ideation	Disjointed flight of ideas
Increased sexual interest	Decreased sexual interest
Anger, verbal aggression	Possible extreme violence
Mild to moderate anorexia	Total anorexia
Inflated self-esteem	Delusions of grandiosity

22.2 Chronic adverse effects

Frequent snorting of cocaine can lead to perforation of the nasal septum (Vilensky, 1982). Reactive hyperaemia of nasal mucosa causes a persistent rhinitis in patients who regularly insufflate cocaine.

Premature atherosclerosis and left ventricular hypertrophy develop in chronic cocaine users. Risks of cocaine use in the short-term are heart attack, respiratory failure, cerebral haemorrhage, renal infarction, seizure, acute state of agitation and overdoses. Cocaine is a vasoconstrictor, which decreases blood flow to the heart and brains temporarily: the risk of heart infarct and stroke increases. People with a weak heart, weak blood vessels, high blood pressure,

diabetes or epilepsy are particularly at risk. Cocaine increases libido, so that one is more inclined to withhold security (e.g. no safe sex).

In high dose, cocaine use can lead to panic attacks or a temporary paranoid psychosis with delusions and hallucinations, i.e. a cocaine delirium, which may subsequently lead to accidents (aggression, violence, suicide). One particularly and frightening type of hallucination is called 'cocaine bugs', which refers to the sensation of tiny creatures crawling over the user's skin. Repetitive scratching (Meyer and Quenzer, 2005) may lead to infection of the wounds. Active cocaine use is the strongest predictor of failure to maintain viral suppression in HIV-positive current and former drug users; 13% of active users maintained suppression versus 46% of non-users (Arnsten et al., 2001).

23 Cocaine-crack

23.1 Acute adverse effects

Problematic drug users in the Netherlands today more often use cocaine in the form of crack or cocaine base than heroin or other opiates. Cocaine is commonly used nasal, but this does not apply to crack cocaine (cocaine base formed from cocaine hydrochloride) which can be smoked. The base coke is smoked with a base pipe ('bases') or foil ('chinezén'; 'chasing the dragon'), and the vapours are inhaled or absorbed through a tube. The vapour inhaled through the base pipe is hotter than that inhaled via 'chinezén' and therefore more harmful to the lungs.

Table 10. Mean physiological and subjective effects of cocaine administered via different routes (Jones, 1990)

Effect	Route (Dose in mg/kg)			
	i.v. (0.6)	Smoked (0.4)	Nasal (3)	Oral (2)
Heart rate increase, BPM (time to max, minutes)	46 (10)	32 (2)	26 (40)	20 (55)
Systolic BP increase mmHg (time to max)	28 (10)	32 (2)	24 (25)	19 (70)
Diastolic BP increase mmHg (time to max)	16 (10)	22 (1)	11 (25)	14 (75)
Pupil diameter increase mm (time to max)	0.8 (4)	1,1 (5)	0,6 (45)	0,5 (90)
Skin temperature decrease °C (time to max)	-2,8 (30)	-1.8 (20)	-4.7 (30)	-5.1 (75)
Subjective high scale (time to max)	48 (4)	35 (1)	18 (20)	18 (70)

23.2 Chronic adverse effects

Crack use is associated to a variety of cardiovascular, respiratory, neurological, and psychiatric problems (Cornish and O'Brien, 1996). The chronic cocaine user tends to administer the drug in high-dose 'binges' interrupted by 'crashes'. Common disorders among heavy crack users are sleep disorders (insomnia followed by exhaustion), eating disorders (appetite suppression alternating with intense hunger) and sexual dysfunction (often impotence). Cocaine (and crack) impairs the immune defence against infections by inhibiting neutrophils and macrophages (Baldwin et al., 1998), so that HIVs more efficiently replicate (Roth et al., 2002, Bagasra and Pomerantz, 1993), leading to a higher viral load and the risk of transmission.

23.3 Disease following chronic cocaine and crack use

Chronic cocaine use is associated with a variety of severe medical conditions. Most complications arise from the cardiovascular toxicity of cocaine (i.e. high blood pressure and irregular heart beat), from the use of contaminated needles, and/or from unsafe sex practices. In particular, the smoking of crack and the intravenous use of cocaine are associated with high-risk sexual practices. In the Netherlands, most (75%) of cocaine-dependent users have problems with other drug use, in particular alcohol (28%), cannabis (20%), heroin (12%), and stimulant drugs (5%) (NDM, 2011). The disease burden of chronic cocaine use, therefore, is difficult to determine. In 2009, the primary diagnosis in hospital-admitted cocaine users (abuse or dependence) was respiratory disease (21%),

accident injury (16%), abuse of alcohol and other drugs (14%), intoxication (8%), cardiovascular disease (8%), and psychosis (4%), respectively. Chronic base cocaine use can lead to over-fatigue and weight loss, leading to complete exhaustion. The physical and mental condition can deteriorate quickly. This depletion in combination with the 'crash' of cocaine users may lead to irritable, sad, aggressive or paranoid conditions.

23.3.1 *Cardiovascular diseases*

Cocaine is a strong vasoconstrictor (within 30 minutes after dosing) which increases the risk of atherosclerosis (Zhou et al., 2004) and coronary disease. Moreover, cocaine alters blood coagulation (platelet activation) (Heesch et al., 2000) and increases the risk of thrombi formation (McKee et al., 2007). Chest pain is a very common symptom in cocaine and crack users. Some feel pain within one hour, others had a delayed onset. Acute myocardial infarction (AMI) is the most commonly reported cardiac consequence of cocaine misuse e.g. (Amin et al., 1990b, Weber et al., 2000). The risk of having an AMI secondary to cocaine use is maximal in the first hour after ingestion, having been reported as 24 times (Mittleman et al., 1999a) or 31 times (Cheng, 2000) the baseline risk. However, lifetime-risk increase has been reported in recent prospective studies to be much lower, with an average risk increase (over non-users) of about 6% (Amin et al., 1990a, Hollander et al., 1995). The incidence of AMI is, however, rather low. In a cohort of 3946 AMI patients, 1% had used cocaine in the last year (Mittleman et al., 1999b). However, more recent studies clearly show a high risk of AMI due to cocaine use in young male adults. Regarding a non-fatal myocardial infarction, the population attributable fraction of cocaine use was 0.9% [Nawrod c.s 2011]. According to Jellinek Clinic one out of four non-fatal heart attacks in men aged <45 years is related to cocaine use. Note that concomitant tobacco smoking is a contributing risk factor. Strokes (CVA) occur in cocaine users (especially in the younger users) (Kaku and Lowenstein, 1990b) and are probably the result of high blood pressure, blood vessel occlusion and vasospasms. Note that in these studies, co-use of amphetamine was often seen (Westover et al., 2007).

23.3.2 *Pulmonary disease*

The lungs are the principal organs exposed to the combustion products of crack cocaine. Crack cocaine (not cocaine i.v.) immediately (Brody et al., 1990) results in airway bronchoconstriction probably via irritation or thermal injury (Tashkin et al., 1996). Cocaine has a wide spectrum of acute pulmonary complications (cf. Table 11) (Thadani, 1996, Smith et al., 1995, O'Donnell et al., 1991, Nadel and Lyons, 1998, Kline and Hirasuna, 1990, Haim et al., 1995a, Ettinger and Albin, 1989, Delaney and Hoffman, 1991). The acute respiratory complaints include cough with sputum production, chest pain with or without shortness of breath, haemoptysis and exacerbation of asthma.

Table 11. Pulmonary complications of smoked cocaine (Haim et al., 1995b)

Acute respiratory symptoms and exacerbation of asthma
Thermal airway injury
Pneumothorax and pneumomediastinum
Pulmonary haemorrhage
Noncardiogenic pulmonary oedema
Pulmonary infiltrates with eosinophilia/interstitial pneumonitis
Pulmonary vascular disease/pulmonary infarction

23.3.3 *Kidney disease*

Chronic cocaine use has been associated with severe nephropathies and renal failure, probably related to infectious diseases, but well-designed prospective epidemiological studies to assess incidence and prevalence of cocaine-associated nephropathies do not exist (Jaffe and Kimmel, 2006a).

23.3.4 *Infections related to needle use*

Epidemiologic research confirms that crack users are at high risk of HIV infection and progression (Kral et al., 1998, Vittinghoff et al., 2001). Crack use is significantly associated with progression to AIDS in HIV-seropositive drug users (Cook et al., 2008, Webber et al., 1999). Compared with non-users, the risk of AIDS-related opportunistic conditions was higher for persistent users and intermittent users during periods of active use, with no difference during periods of abstinence (Lucas et al., 2006). Risk of HIV transmission or re-infection among HIV-positive crack users may also be increased because they are more likely to delay or reduce health care utilization (Kang et al., 2006, Cunningham et al., 2006), they poorly comply to antiretroviral medications (Moss et al., 2004, Hinkin et al., 2007), and they may experience increased viral load and greater risk of transmission. It has been suggested that there is a connection between binge use and transmission risk. Most crack smokers use in binge cycles; that is, users rarely stop with one hit, but use as much crack as resources allow, and then "crash" into a state of physical and psychological withdrawal (Harzke et al., 2009a).

HIV-positive persons who use crack cocaine are frequently sexually active and seem to be at higher risk for HIV transmission or re-infection (Harzke et al., 2009b, Campsmith et al., 2000). The reason may be the high prevalence of unprotected sex, having multiple partners, and exchange sex for drugs or money.

Skin and soft tissue infections are common among injecting drug users. Injection of cocaine with non-sterile needles gives abscesses at the injection site and infections, like viral hepatitis and AIDS. In several studies, prevalence rates of 20-30% have been reported in Europe and 16-65% in the United States (Ruiz and Strain, 2011). In a prospective study among drug injectors in Amsterdam, the incidence of abscess was 33 per 100 person-years. Skin and soft tissue infections may progress to systemic infections including endocarditis. Finally, crack cocaine use is associated with tuberculosis (Story et al., 2008).

23.4 **Mortality**

Death caused by cocaine overdose is low in the Netherlands, about 23 cases per year during 2000-2009, and is highest in middle-age users (aged 30-44 years). In 2008, the EU-15 reported about 1000 cocaine-related deaths (EMCDDA, 2010). Most fatalities associated with cocaine use are caused by cardiovascular or cerebrovascular accidents. Therefore, chronic cocaine use may be overlooked as the cause of death and reported death rates likely underestimate the real death rate by cocaine.

23.5 **Economic burden**

The health burden related to cocaine use in Europe is not yet identified and probably increasing (EMCDDA, 2010). Approximately 5% to 10% of emergency department visits in the United States is believed to be secondary to cocaine usage, leading to the evaluation of approximately 64,000 patients annually for possible myocardial infarction, of which approximately 57% are admitted to the hospital, resulting in an annual cost exceeding € 60 million (Maraj et al., 2010a).

According to Maraj et al. (Maraj et al., 2010b), there is no evidence to suggest that pre-existing vascular disease is a prerequisite for the development of a cocaine-related cardiovascular event.

24 Alcohol

24.1 Acute adverse effects

The immediate negative effects of alcohol on the circulation are relatively minor and reversible. Overindulgence will produce early symptoms of alcohol intoxication, like nausea and vomiting, which urges most people to temporarily stop drinking alcohol. Headache (hang-over), impaired sexual capability and temporary loss of memory are typical symptoms of high alcohol consumption and intoxication. The consumption of high amounts of alcohol (blood alcohol concentrations >400 mg/dl; 4‰) produces loss of consciousness, and can be dangerous or even prove lethal due to respiratory depression and coma. Moreover, due to the diminished reaction time and impaired locomotor activity, alcohol intoxicated subjects are more liable to sometimes fatal traffic accidents and injuries. Alcohol overconsumption is also associated with a variety of risk-taking behaviours, which may result in accidents and injuries (drowning, chronic disability, HIV-infection in the case of unprotected sexual activity) (WHO, 2006). In alcoholics, the risk of death by suicide, homicide, fire, and drowning is roughly doubled. In Europe, alcohol is involved in 40% of murders and manslaughters, and in 16% of suicides (Anderson and Baumberg, 2006).

24.2 Disease due to chronic excessive alcohol use

24.2.1 Introduction

Alcohol and tobacco are legal drugs, but just because they are legal does not imply that these drugs should be considered safe. World-wide, approximately 125 million people are affected by alcohol-use disorders and many more people suffer from alcohol use disorders than from illicit drug use disorders. Annually, alcohol kills 35 people per every 100,000, whereas for illicit drugs this is nine times less. In 2004, 7.6% of all global burdens of disease and injuries among men, and 1.4% among women, were linked to alcohol use.

Table 12. Proportions attributable to alcohol use for major alcohol related diseases and injuries (Room et al., 2005)

	Men	Women	Both
Mouth and oropharynx cancers	22%	9%	19%
Oesophageal cancer	37%	15%	29%
Liver cancer	30%	13%	25%
Breast cancer	-	7%	7%
Unipolar depressive disorders	3%	1%	2%
Epilepsy	23%	12%	18%
Ischaemic heart disease	4%	-1%	2%
Haemorrhagic stroke	18%	1%	10%
Ischaemic stroke	3%	-6%	-1%
Liver cirrhosis	39%	18%	32%
Traffic accidents	25%	8%	20%
Drownings	12%	6%	10%
Homocide	26%	16%	24%

Alcohol and tobacco have a high prevalence (75% and some 20-30%, respectively), whereas the prevalence of illicit drug use is much lower, i.e. up to 4%, in Western countries (WHO, 2010). This explains why the population level social harm scores for legal drugs are generally higher than individual level social harm scores (and thus the total harm score), whereas the opposite is generally true for illicit drugs (cf. Table 1).

24.3 General protective and harmful effects

According to the latest estimates of the WHO (WHO, 2010, WHO, 2011), the global burden of disease attributed to alcohol and illicit drugs amounts to 5.4% of the total burden of disease. The finding that high alcohol consumption is potentially harmful is also corroborated by the fact that it is the third main cause of early death and illness in the EU with 4.5 million DALYs lost and only more DALYs lost due to tobacco use (7.5 million) and hypertension (6 million) (Anderson and Baumberg, 2006). In 2006, the epidemiological studies on long-term effects of alcohol consumption have been reviewed by the Dutch Health Council (Health Council of the Netherlands, 2006) and parts are used in the present review.

Table 13. Relative risks (RRs) for selected medical conditions of alcohol consumption (10-30 g per day; 2-3 drinks per day) by men and women in three age categories [33]

Medical condition	Relative risk					
	Women (age in yr)			Men (age in yr)		
	0-19	20-39	40+	0-39	40-59	60+
Cirrhosis of the liver	1.3	9.5	13.0	1.3	9.1	13.0
Acute and chronic pancreatitis	1.3	1.8	1.8	1.3	1.8	3.2
Epilepsy	1.3	7.2	7.5	1.2	7.5	6.8
Mouth and oropharynx cancers	1.5	2.0	5.4	1.5	1.9	5.4
Oesophageal cancer	1.8	2.4	4.4	1.8	2.4	4.4
Laryngeal cancer	1.8	3.9	4.9	1.8	3.9	4.9
Liver cancer	1.5	3.0	3.6	1.5	3.0	3.6
Breast cancer	1.1	1.4	1.6			
Other neoplasms	1.1	1.3	1.7	1.1	1.3	1.7
Hypertension	1.4	2.0	2.0	1.4	2.0	4.1
Coronary heart disease	0.8	0.8	1.1	0.8	0.8	1.0
Ischaemic stroke	0.5	0.6	1.1	0.9	1.3	1.7
Haemorrhagic stroke	0.6	0.7	0.8	1.3	2.2	2.4
Cardiac arrhythmias	1.5	2.2	2.2	1.5	2.2	2.2
Spontaneous abortion	1.2	1.8	1.8			
Low birth weight ¹	1.0	1.4	1.4	1.0	1.4	1.4
Prematurity ¹	0.9	1.4	1.4	0.9	1.4	1.4
Intra-uterine growth retardation ¹	1.0	1.7	1.7	1.0	1.7	1.7

¹Relative risk refers to drinking of mother.

Moderate alcohol consumption has a protective effect on cardiovascular disease; regular consumption of small amounts of alcohol being more protective than the same amount taken in larger doses less frequently (Grobbee et al., 1999). The few cohort studies conducted among women all confirm the protective effects of alcohol at consumption levels below a rather high (see below) cut-off level of 48 g/day (Fuchs et al., 1995, Klatsky et al., 1997, Rehm et al., 1997, Thun et al., 1997b). It is accepted that drinking alcohol confers a significant health benefit in terms of reduced coronary heart disease (CHD) mortality and morbidity on men aged over 40 and postmenopausal women, and a maximum oral intake of 24 g and 32 g of alcohol per day for women and men, respectively is recommended (UKDH, 1995). However, it has been suggested that the cardiac protection caused by alcohol is overestimated in prospective epidemiological mortality studies, because of contamination of the abstainer category with occasional or former drinkers (Filmore et al., 2006). The WHO acknowledges the evidence of a protective effect only at levels of drinking as low as one drink per week, but advised that any beneficial health effects be sought by other means because of the many adverse effects of alcohol (WHO, 2001). Indeed, the harm of alcohol consumption is many times higher than its protective effects, considering that consumption of alcohol is related to over 60 medical conditions. This is confirmed by the data depicted in Table 12, which reflect the proportions attributable to excessive alcohol use for major alcohol related diseases and injuries.

Based on epidemiological data, the UK Interdepartmental Working Group on Sensible drinking (UKDH, 1995) confirmed the J-shaped relationship between alcohol consumption and all-cause mortality. Non-drinkers have higher all-cause mortality than light and moderate drinkers, and heavy drinkers have even higher all-cause mortality than either group. Table 13 depicts the relative risks (RRs) for selected medical conditions of alcohol consumption.

A meta-analysis, published in 1996, of 16 cohort studies on alcohol consumption and all-cause mortality confirms the J-shape curve (Holman et al., 1996c). At consumption levels considered as 'responsible' (i.e. <20 g/day for women and <40 g/day for men), the relative risks for alcohol-related cancers and liver cirrhosis compared to abstainers were significantly increased, whereas those for ischaemic heart disease, stroke and cholelithiasis were significantly decreased (Holman et al., 1996b). An important prospective cohort study not yet included in the latter review nor in the meta-analysis, was the study by Thun et al. among 490,000 US men and women (mean age 56 years; range 30 to 104), who reported their alcohol and tobacco use in 1982 (Meister et al., 2000, Thun et al., 1997a). A nine-year follow-up of these individuals showed that 46,000 of them died. The analyses were adjusted for many potential confounders, including education and smoking. For men as well as women, total mortality appeared to be lowest at a consumption of one alcoholic drink per day (12 g/day), but in women the rate of increase of the risk at a higher consumption level was larger than in men.

24.3.1 *Disease in the digestive tract (non-carcinogenic)*

High alcohol consumption frequently leads to gastritis, ulcers and severe stomach bleedings. In addition, alcohol induces lesions of the oesophagus and duodenum and is also an aetiological factor in acute and chronic pancreatitis (Rall, 1992). In humans, the most critical and dominant non-carcinogenic effect induced by alcohol overconsumption appears to be liver cirrhosis.

Liver cirrhosis caused by excessive drinking is increasing rapidly, with a tenfold increase in UK in the last 30 years (Anderson and Baumberg, 2006). Though only few prospective cohort studies on alcohol consumption and liver cirrhosis

have been conducted, there is no doubt that excessive alcohol use often results in liver damage and ultimately death due to liver cirrhosis. Reversible conditions, such as steatosis (fatty liver) and alcoholic hepatitis, precede the occurrence of irreversible cirrhosis and are presumably causally related to it (Sorensen et al., 1984). The magnitude of the relative risks (RRs) of three large cohort studies appeared to be quite comparable (Becker et al., 2002, Holman et al., 1996a, Klatsky, 1994). A consumption of 12 g of alcohol per day did not seem to increase the risk of liver cirrhosis to a very large extent, but above that level, an increase of the risk was apparent. Typically, cirrhosis requires the consumption of at least 80 g of alcohol daily for 10-20 years (Lelbach, 1975). It was not clear, however, whether women are at a higher risk than men.

In individuals who present with a long history of gastroesophageal reflux disease, there is an increased incidence of Barrett's oesophagus. Barrett's oesophagus, a metaplastic conversion of the lining of the lower oesophagus, is a known precursor lesion for oesophageal cancer (Johnson and Marzani-Nissen, 2010).

Pancreatitis, both acute and chronic, is another complication of excessive alcohol consumption; it ranges from an uneasy but stable condition to a medical emergency, depending on the severity of the event.

Both type I and type II diabetes may be caused by excessive alcohol consumption. The development of type I diabetes is rare and due to almost complete destruction of the pancreas. Type II diabetes is more common due to weight gain and carbohydrate intake. Hypogonadism and osteoporosis are other complications. Thyroid disease can also result in excessive alcohol consumption, drug abuse or dependence (Johnson and Marzani-Nissen, 2010). Alcohol can also exacerbate hepatitis C infection, considering that more than half of all patients with hepatitis C have a past history of alcohol use, and chronic alcohol consumption (Safdar and Schiff, 2004). Individuals that consume more than five drinks per day with hepatitis C show an increase in the rate of liver fibrosis, cirrhosis, hepatocellular carcinoma and, possibly, death from liver disease (Jamal et al., 2005).

Gout is another common complication of chronic, excessive alcohol consumption. Podagra (pain in the big toe) is a typical complaint. The use of alcohol also seems to mitigate certain autoimmune diseases, like systemic lupus erythematosus and rheumatoid arthritis.

24.3.2 *Cardiovascular disease*

Focused on cardiovascular disease, the International Life Sciences Institute (ILSI) report reviewed the available scientific and epidemiological evidence from case-control and cohort studies for the association of alcohol consumption and conditions like coronary heart disease, lipids, haemostasis, atherosclerosis, blood pressure, insulin sensitivity, and (different types) of stroke (Grobbee et al., 1999). A remarkable consistency across diverse populations was observed in the studies with coronary heart disease (CHD) mortality or incident CHD (or myocardial infarction) as the endpoint (Grobbee et al., 1999). The authors observed a U-shape relationship between alcohol dose and CHD. The level at which fatal and non-fatal CHD signs begin to increase could not be clearly identified from the U-shaped curve but appeared to be somewhere between two and six drinks/day (20-60 g/day), a range that is hardly useful for prevention strategies and policy making (Grobbee et al., 1999).

Whatever the cause, the incidence of cardiac arrhythmia doubles for heavy drinkers compared with light drinkers (Cohen et al., 1988). The incidence of cardiac arrhythmias following alcohol consumption is commonly known as 'holiday heart phenomenon'. This follows the observation that supraventricular

arrhythmias are most common in alcoholics on Mondays and between Christmas and New Year (Ettinger et al., 1978, Menz et al., 1996). Alcohol use leads to hypertension as a causal relation exists between the use of >30-60 g/day and blood pressure elevation in men and women (Grobbee et al., 1999). Assuming a linear relationship with no threshold, an additional drink a day (10 g) would increase both systolic and diastolic blood pressures by 1-2 mmHg (Anderson et al., 1993a, UKDH, 1995). A generally accepted clinical view would be that for men the rise in blood pressure produced by 32 g/day (about 6 mmHg systolic blood pressure and 4 mmHg diastolic) would give rise for concern (UKDH, 1995). Though most epidemiological studies suggest that regular light to moderate alcohol intake (16-32 g/day) probably reduces the risk of ischaemic stroke, regular consumption of more than 40 g of alcohol per day and binge drinking increases the risk of ischaemic and haemorrhagic stroke (due to cerebral or subarachnoid haemorrhage) (Anderson et al., 1993b, Grobbee et al., 1999, UKDH, 1995).

24.3.3 *Cancer disease*

Numerous epidemiological studies have shown a causal relation between alcohol consumption and cancers of the oral cavity, pharynx, larynx, oesophagus, liver, colon, rectum and female breast cancer (Baan et al., 2007, Rehm et al., 2010c) (for relative risks cf. Table 3). Renal cell cancer, and non-Hodgkin's lymphoma show much weaker, and less consistent associations with alcohol consumption (Baan et al., 2007, Corrao et al., 1999, Doll, 1998, Longnecker, 1995, Rehm et al., 2010b). A causal relation between alcohol and colorectal cancer was established only recently by the International Agency for Research on Cancer (IARC) (Baan et al., 2007, Rehm et al., 2010a). A meta-analysis of 27 studies (follow-up and case-control studies) (Longnecker et al., 1990) showed a RR for colon cancer of only 1.10 (CI: 1.05-1.14) for two drinks per day (equivalent to approximately 24 g of alcohol per day). Later reports showed similar effect sizes, though at higher daily alcohol consumption (Cho et al., 2004a, Corrao et al., 2004, Moskal et al., 2007, Rehm et al., 2010d). As an example, the analysis of eight pooled cohort studies, which included almost half a million subjects and 4,687 cases of colorectal cancer, gave a RR of 1.16 (CI: 0.99 - 1.36) for persons who consumed 30 to less than 45 g/day and 1.41 (CI: 1.16 - 1.72) for those who consumed 45 g/d or greater compared with non-drinkers (Cho et al., 2004b).

A statistically significant, but weak association between alcohol consumption and breast cancer was observed in women by a meta-analysis of six pooled prospective cohort studies: RR of 1.09 (CI: 1.04 - 1.13) for each 10 g of alcohol up to 60 g per day appeared to be linearly related to alcohol consumption (Smith-Warner et al., 1998). Hamajima et al. (Hamajima et al., 2002) pooled the data of 58,515 breast cancer cases and 95,067 controls from 53 cohort and case-control studies, which represented 80% of all data available on this subject world-wide. They also observed an almost linear dose-response association, but estimated a (confounder-adjusted) slightly lower increased risk of 7.1% overall for every 10 g of alcohol consumption. A meta-analysis of five US cohort studies published since 1990 yielded a similar low relative risk of 1.06 (95% CI: 1.00-1.11) for consumers of 12 g per day, as compared with non-drinkers (Ellison et al., 2001).

Moreover, heavy alcohol consumers are often also regular smokers. As such, it has been suggested that alcohol may act as co-carcinogen by enhancing the carcinogenic effects of tobacco smoking (Blot et al., 1988).

24.3.4 *Hematologic/hematopoietic system*

Anaemia, both macrocytic and microcytic, are possible. Macrocytic anaemia due to folic acid or vitamin B₁₂ deficiency. Note that an elevated mean corpuscular volume may also result from liver disease in the lipid bilayers that red cells do not form properly. When liver disease is severe, platelets may be destroyed or can isolate an enlarged spleen. Microcytic anaemias are associated with active bleeding or bleeding and the evaluation should ask for a gastrointestinal disorder or injury. Sideroblastic anaemia may also occur (Johnson and Marzani-Nissen, 2010).

24.3.5 *Other diseases*

Psoriasis vulgaris, acne rosacea, and erythropoietic protoporphyria are all common skin conditions associated with excessive alcohol consumption. With liver disease, spider nevi, telangiectasias, palmar erythema (red palms), spider angiomas, and hepatic porphyria, especially porphyria cutanea tarda (bullous erosions, blistering, crusting lesions and scars heal with hyperpigmentation or depigmentation on the face, side of the neck and the backs of the hands) could be found (Johnson and Marzani-Nissen, 2010). Vitamin and essential nutritional deficiencies due to poor food intake or impaired gastrointestinal and hepatic function of the alcoholic seem to cause many psychiatric syndromes that are common in alcoholics, such as Wernicke's encephalopathy, Korsakoff's psychosis, polyneuritis, and encephalopathy (Rall, 1992). The most important of these and commonly seen in alcoholics is Wernicke-Korsakoff syndrome. Alcohol is immunosuppressive so that its consumption increases the susceptibility to infectious diseases (pneumonia, tuberculosis, hepatitis C and HIV) (NIAAA, 2000).

24.4 **Mortality**

Alcohol consumption accounts for 85,000 deaths per year in the U.S., tobacco smoking for a tenfold higher rate, whereas illegal drugs account for 'only' for one-fifth of the alcohol-related deaths (17,000). Worldwide, alcohol causes 1.8 million deaths per annum. Alcohol causes a considerable disease burden: 3.2% of the global deaths and 4.0% of the global DALYs in the year 2000 could be attributed to this exposure (Rehm et al., 2003, WHO, 2006). Alcohol consumption is related to more mortality in young people than it is in any other age group. In European countries, over 10% of female mortality and 25% of male mortality in those aged 15-29 years is alcohol-related. In the EU, alcohol consumption at harmful levels leads to 195,000 deaths each year (various cancers, liver cirrhosis, road traffic and other accidents, homicides, suicides and neuropsychiatric conditions) (Anderson and Baumberg, 2006, EUPHIX, 2007, Rehm et al., 2006b). Taking into account the prevention of deaths by moderate consumption, alcohol causes an estimated 115,000 deaths in people up to the age of 70 each year in the 25 countries of the EU (Anderson and Baumberg, 2006). In Europe, a rise of one litre per capita in alcohol intake was associated with a 1% - rise in all causes of morbidity (Her and Rehm, 1998).

24.5 **Financial burden related to alcohol consumption**

The costs of 'social harm' due to excessive alcohol consumption outweigh the health costs. Alcohol consumption also has a large economic benefit (employment, tax duties). Both items are, however, no topic of the present review.

The total alcohol-related costs in the EU average is about 270 billion euro per year (Anderson and Baumberg, 2006), and the total health impact attributed to

alcohol consumption (due to alcohol disorder and the secondary health effects linked to alcohol consumption) for all three EU regions is estimated to be 6.1% of all deaths and 10.7% of all DALYs. An estimated 11.9% of all DALYs in men and 1.4% of all DALY in women can be attributed to the total disease burden of alcohol in the western EU member states (Eastern Europe and former Soviet states excluded) (Rehm et al., 2006a), which does not include the social harm experienced by family members or by victims of crime and accidents (EUPHIX, 2007).

Based on a review of existing studies, the total tangible, i.e. verifiable, cost of alcohol to EU society in 2003 was estimated to be 125 billion euro, equivalent to 1.3% of the Gross Domestic Product (GDP) (Anderson and Baumberg, 2006, ICAP, 2006). The intangible costs, related to the value people place on pain, suffering and lost life that occurs to the criminal, social and health harms caused by alcohol, are estimated to be € 270 billion or 2.8% of the GDP (Anderson and Baumberg, 2006, ICAP, 2006).

Balakrishnan et al. (Balakrishnan et al., 2009) estimated that alcohol consumption was responsible for 31,000 deaths in the UK in 2005 and that the consumption of alcohol costs the UK National Health Service (NHS) £ 3.0 billion in 2005-06. Alcohol consumption was responsible for 10% of all disability adjusted life years in 2002 (male: 15%; female: 4%) in the UK.

Table 14. Social costs of excessive alcohol consumption EUR per year (KPMG 2001b)

Type of costs	Amount (euro)
Care and treatment of addiction	68 million
General healthcare	115 million
Work	1,554 million
Crimes and offences	840 million
Total	2,577 million

Finally, alcohol is often used in combination with other drugs. For instance, we recently showed that the consumption of magic mushrooms is relatively safe, but can occasionally lead to fatal accidents when used in combination with alcohol (Van Amsterdam et al., 2011).

The treatment of alcohol addiction is much higher than that of illegal drugs because the number of alcoholics in Netherlands is 13 times higher than the number of illicit drug addicts, and alcohol- and tobacco-related deaths are 15 and 333 times higher respectively than those related to the use of illicit drugs (Council, 1999, Trimbos, 2004). In the Netherlands, the financial burden of addiction treatment (alcohol, drugs, and gambling) amounts to € 197 million, of which 34 per cent was attributable to alcohol (KPMG, 2001). The costs of hospitalisations related to excessive alcohol use amount to € 106 million.

25 Tobacco

25.1 Acute adverse effects

Immediately after exposure to nicotine, a rush of adrenaline stimulates the body and causes a sudden release of glucose, as well as an increase in blood pressure, respiration, and heart rate (NIDA, 2006). Nicotine also suppresses insulin output from the pancreas, which means that smokers are always slightly hyperglycemic (NIDA, 2006). Other negative effects, including nausea and dizziness, rapidly disappear during the everyday cycle of use (INSERM, 2004). Smoking a single cigarette decreases the cutaneous blood flow in habitual smoker as well as in non-smoker subjects (Monfrecola et al., 1998). In naïve smokers, the irritating smoke induces cough. The well-known smokers' cough, on the other hand, happens since their cilia are so damaged that phlegm can only be removed by coughing (USDHHS, 2006).

25.2 Diseases

The annual figures for hospital admissions as a direct result of smoking in the Netherlands were estimated in 2005. About 90,000 people of 35 years and older were admitted to hospital with a smoking related disease (Cruts et al., 2008). This was about 7.5 % of the total hospital admissions within this group. Mainly airway related cancer, cardiovascular disease and chronic airway obstruction were the most frequent reasons for admission. Acute intoxications by tobacco were primarily recorded in young children (365 in 2009), as a result of nicotine poisoning through the accidental swallowing of a cigarette or rolling tobacco (NDM, 2011).

Table 15. Diseases and causes of death related to smoking (NDM, 2011)

Disease	Total deaths		Smoking related deaths	
	Male	Female	Male	Female
Lung cancer	6,427	3,533	5,885 (91.6%)	2,585 (73.2%)
COPD	3,510	2,741	2,966 (84.5%)	1,885 (68.8%)
Coronary disease	6,183	4,481	1,710 (27.7%)	549 (12.3%)
Stroke	3,453	5,602	684 (19.8%)	541 (9.7%)
Heart failure	2,550	4,004	435 (17.1%)	204 (8.0%)
Oesophageal cancer	1,149	420	913 (79.5%)	262 (62.4%)
Laryngeal cancer	162	38	130 (80.3%)	32 (84.2%)
Apertural cancer	406	216	374 (92.1%)	118 (54.6%)
Total	23,840	21,035	13,069 (54.8%)	6,176 (29.4%)

Contribution of smoking related deaths are given in percentages between brackets.

25.2.1 Cancer disease

The overall rates of death from cancer are two to four times as high among smokers as non-smokers (NIDA, 2006). The risk of dying from lung cancer is 13 times (women) and 23 times (men) higher compared with never smokers (USDHHS, 2006). Smoking causes about 90% of lung cancer deaths in men and almost 80% of lung cancer deaths in women (NIDA, 2006, WHO IARC, 2004), which equals 8,000 annual deaths in The Netherlands (RIVM, 2008). Tobacco smoking increases the risk of all histological types of lung cancer including squamous-cell carcinoma, small-cell carcinoma, adenocarcinoma (including bronchiolar/-alveolar carcinoma) and large-cell carcinoma (WHO IARC, 2004).

Smoking is also associated with cancers of the bladder, ureter, oral cavity, pharynx, larynx (voice box), oesophagus, cervix, kidney, pancreas, the nasal cavities and nasal sinuses, liver, stomach, and acute myeloid leukaemia. (NIDA, 2006, USDHHS, 2006, WHO IARC, 2004), risks being two- to threefold higher than in non-smokers (WHO IARC, 2004) and amounting to 1,700 deaths in total in The Netherlands (RIVM, 2008).

25.2.2 *Respiratory Disease*

Smoking causes chronic obstructive lung disease (COPD) and exacerbates asthma symptoms (Hylkema et al., 2007, NIDA, 2006, USDHHS, 2006). Cigarette smoking is associated with a tenfold increase in the risk of dying from COPD (Hatsukami et al., 2008). More than 90 per cent of all deaths from COPD is attributable to cigarette smoking (NIDA, 2006, USDHHS, 2006). The biological processes leading to the development of COPD are oxidative stress and pulmonary inflammation (USDHHS, 2006, Wilford et al., 2006). Other smoking related respiratory symptoms are coughing, phlegm, wheezing, dyspnoea and impaired lung growth (USDHHS, 2006).

25.2.3 *Cardiovascular disease*

Smoking substantially increases the risk of heart disease, including acute myocardial infarction, sudden cardiac death, stroke, aortic aneurysm, and peripheral vascular diseases (Bullen, 2008a, NIDA, 2006, Rahman and Laher, 2007b). Cigarette smokers are two to four times more likely to develop coronary heart disease than non-smokers, more than ten times as likely to develop peripheral vascular disease, and death from rupture of an atherosclerotic abdominal aneurysm is also more common in smokers (USDHHS, 2006). Tobacco smoke is both prothrombotic and atherogenic (Bullen, 2008b, Rahman and Laher, 2007a) by narrowing the blood vessels (atherosclerosis) (Leone, 2007, Rahman and Laher, 2007c).

25.2.4 *Other diseases*

Tobacco is an important risk factor for periodontal diseases (Jacob et al., 2007, Kinane and Chestnutt, 2000). With increased use of tobacco, patients show higher periodontal probing depths, increased clinical attachment loss, more alveolar bone resorption, a higher prevalence of gingival recessions, and a higher risk for tooth loss (Jacob et al., 2007, Saxer et al., 2007). Cigarette smoking has a negative effect on wound healing and periodontal treatment procedures (Jacob et al., 2007). Women who smoke have an increased risk for cataract, hip fractures, low bone density, and peptic ulcer disease when *Helicobacter pylori* positive (Hatsukami et al., 2008, USDHHS, 2006).

25.3 **Environmental tobacco smoke**

Environmental tobacco smoke (ETS) is a major source of indoor air contaminants and causes disease in non-smoking subjects. The Dutch Health Council (Gezondheidsraad) estimates that ETS causes approximately a few hundred lung cancer deaths, a few thousand cardiovascular deaths, and many tens of thousands of cases of respiratory diseases with children (Gezondheidsraad, 2008). Never-smokers exposed to ETS have a statistically significant increase in risk of some 10-30% to develop lung disease, and the risk of an acute coronary heart disease event is increased by some 25-35% (USDHHS, 2006, WHO IARC, 2004). ETS further causes acute respiratory infections and hearing problems (USDHHS, 2006), and is especially in foetuses

and children, associated with poorer neurocognitive performance (Swan and Lessov-Schlaggar, 2007).

25.4 Mortality

Smoking has been estimated to be the main cause of premature mortality in the Netherlands. In 2009 there were 19,245 deaths of people 20 years and older attributed to smoking (NDM, 2011). This number has remained relatively constant throughout previous years. These numbers are surely an underestimate, since the effects of passive smoking, through environmental smoke, have not been included and are difficult to exactly estimate. These number of deaths through passive smoking were estimated to be at least several thousands in 2009. Worldwide, the incidence of death through passive smoking is estimated to be one on every hundred deaths (Oberg et al., 2011). Main causes of death include cardiovascular disease, airway infections and lung cancer.

25.5 Economic burden

Globally, smoking has been estimated to cause 5 million deaths per year, and if present trends continue, 10 million smokers per year are projected to die by 2025 (Davis et al., 2007, Hatsukami et al., 2008). In the Netherlands, smoking caused almost 20,000 deaths in 2006, of which some 8,000 were due to lung cancer, 4,900 due to COPD, 4,700 due to CVD, and 1,700 due to other types of cancer (RIVM, 2008). The economic health care burden of smoking in the Netherlands were investigated in 2003; it was estimated that around € 2 billion were spent on health care attributed to the chronic effects of smoking; 3.7% of the total amount was spent on health care (van Baal et al., 2006). This equalled € 104 per inhabitant in the Netherlands per year. This is a higher amount than is spent on health care resulting from overweight/obesity. Surprisingly, in the long run, smokers save on governmental spending in comparison with healthy non-smokers, mainly because their lifespan expectancy is shortened (van Baal et al., 2008). In this way, it was estimated that smokers save up € 60,000 on lifetime health care in comparison with non-smokers. It was shown that health care costs of men who do not smoke are 15% more than those of smokers. For women the difference was slightly higher with 18% (Barendrecht et al., 1997). Smoking cost the National Health Service (NHS) about € 1.6 billion in 1991 and over the past 10 years has been responsible for about 100,000 deaths yearly (Allender et al., 2009a). Allender et al. (Allender et al., 2009b) estimated that the deaths attributable to smoking in 2005 was 19% of all deaths (27% in men and 11% in women). Smoking was directly responsible for 12% of disability adjusted life years lost in 2002 (15.4% in men; 8.5% in women) and the direct cost to the NHS was £ 5.2 billion in 2005-6.

Economically, more than € 55 billion of total U.S. healthcare costs each year are attributable directly to smoking, excluding medical care costs associated with disease caused by second-hand smoke (NIDA, 2006). The costs related with absenteeism from work (USDHHS, 2006) and the economical profit from dying at an earlier age (reduction in old age pension and ageing-related health care costs) and excise duties are no topic of this review.

26 Acknowledgements

The present study was financially supported by The Netherlands Organisation for Health Research and Development (ZonMw). The authors are especially thankful to Drs. Bert Molendijk of The Centre for Special Dentistry of Jellinek for his contribution about the adverse oral health effects in drug users.

Our special thanks goes to the experts Roel Kerssemakers, Gabe Sonke, Jan Krul, Irma de Vries, Armand Girbes, Barbara Broers, Ton Nabben, Dirk Korf, Magda Boonstra, Margriet van Laar, Ed Pennings, Raymond Niesink, Evelien Vermeulen en Hein Sigling, who have advised us in the identification of the gaps in knowledge.

27 Accountability

We examined the identified peer-reviewed literature on human studies in PubMed from January 1990 to November 2011 (approximately the last 20 years) in English, German, Dutch and French language. The search was not confined to the population-based controlled trials or prospective studies in order to retrieve also the information about drug abuse and physical illness from other studies (e.g. case studies, single blind studies). Because the focus of the search and subsequent review of the literature was on humans, animal studies in connection with illegal substances abuse were not included in the search. A broad search strategy was conducted to ensure retrieval of all data about the physical illnesses of the 19 drugs selected. Finally, the experts engaged were asked to report unpublished data to the authors.

PubMed literature search strategy

Mesh terms used: 'Diseases category/aetiology' of 'Diseases

category/epidemiology', 'adverse effects', 'Comorbidity', 'Comorbid'.

Free text terms: the names of the 19 drugs and their synonyms (e.g. marihuana OR cannabis OR hemp OR hashish).

The Medical Subject Headings (MeSH) terms were combined with the free text terms of the 19 drugs.

Textbooks in addiction medicine and internal medicine do not adequately review the physical morbidity associated with drugs of abuse. Nevertheless, the following seven textbooks were used as a source of physical health problems associated with drugs of abuse.

1. Bankole A. Johnson. *Addiction Medicine: Science and Practice*. Springer, 2010.
2. Pedro Ruiz and Eric C. Stra. *Lowinson and Ruiz's Substance Abuse: A Comprehensive Textbook*. 5th Ed. Lippincott Williams & Wilkins, 2011.
3. Adam J. Gordon. *Physical Illness and Drugs of Abuse. A Review of the Evidence*. Cambridge University Press. 2011.
- 4a. N.L. Benowitz, Sh. Zevin Chapter 8. *Medical Aspects of Drug Abuse*. In: *Addiction and the medical complications of drug abuse*. Zevin and Benowitz. Ed. S.B. Karch. CRC Press 2007.
- 5b. B.A. Roth, N.L. Benowitz and K.R. Olson. B.A. Roth. Chapter 9. *Emergency Management of Drug Abuse*. In: *Addiction and the medical complications of drug abuse*. Zevin and Benowitz. Ed. S.B. Karch. CRC Press 2007.
6. M.A. Schuckit. *Drug and alcohol abuse: a clinical guide to diagnosis and treatment*. 6th Ed. Springer 2006.
7. A.J. Gordon. *Physical Illness and Drugs of Abuse: A review of the evidence* Cambridge University Press. 2010.

References found in the different retrieved sources have been used as further source of information.

This paper is to rather be regarded as a 'bibliography' than a critical review, because the aim of this paper is to summarize data from literature about physical harm of drug abuse/addiction.

28 References

- Abel EL: Marijuana and sex: a critical survey. *Drug Alcohol Depend* 1981; 8(1):1-22.
- Abraham DH. Hallucinogen-related disorders. In B.J. Sadock & V.A. Sadock (Eds), *Kaplan & Sadock's Comprehensive Textbook of Psychiatry* (8th ed.). Philadelphia, PA: Lippincott, Williams & Wilkins, pp. 1015-1024. 2004.
- Agaba EA, RM Lynch, A Baskaran, T Jackson: Massive intracerebral hematoma and extradural hematoma in amphetamine abuse. *Am J Emerg Med* 2002; 20(1):55-57.
- Al Hadrani AM: Khat induced haemorrhoidal disease in Yemen. *Saudi Med J* 2000; 21(5):475-477.
- Al Motarreb A, M Al Kebsi, B Al Adhi, KJ Broadley: Khat chewing and acute myocardial infarction. *Heart* 2002; 87(3):279-280.
- Al-Hebshi NN, N Skaug: Effect of khat chewing on 14 selected periodontal bacteria in sub- and supragingival plaque of a young male population. *Oral Microbiol Immunol* 2005; 20(3):141-146.
- Al-Motarreb A, S Briancon, N Al-Jaber, B Al-Adhi, F Al-Jailani, MS Salek, KJ Broadley: Khat chewing is a risk factor for acute myocardial infarction: a case-control study. *Br J Clin Pharmacol* 2005; 59:574-581.
- Aldington S, M Harwood, B Cox, M Weatherall, L Beckert, A Hansell, A Pritchard, G Robinson, R Beasley: Cannabis use and risk of lung cancer: a case-control study. *Eur Respir J* 2008; 31(2):280-286.
- Ali AA, AK Al Sharabi, JM Aguirre, R Nahas: A study of 342 oral keratotic white lesions induced by qat chewing among 2500 Yemeni. *J Oral Pathol Med* 2004; 33(6):368-372.
- Ali WM, M Zubaid, A Al-Motarreb, R Singh, SZ Al-Shereiqli, A Shehab, W Rashed, NQ Al-Sagheer, AH Saleh, J Al Suwaidi: Association of khat chewing with increased risk of stroke and death in patients presenting with acute coronary syndrome. *Mayo Clin Proc* 2010; 85(11):974-980.
- Allender S, R Balakrishnan, P Scarborough, P Webster, M Rayner: The burden of smoking-related ill health in the UK. *Tob Control* 2009; 18(4):262-267.
- Amin M, G Gabelman, J Karpel, P Buttrick: Acute myocardial infarction and chest pain syndromes after cocaine use. *Am J Cardiol* 1990a; 66(20):1434-1437.
- Amin M, G Gabelman, J Karpel, P Buttrick: Acute myocardial infarction and chest pain syndromes after cocaine use. *Am J Cardiol* 1990b; 66:1434-1437.
- Anderson K: Bronchospasm and intravenous street heroin. *Lancet* 1986; 1(8491):1208.
- Anderson P, B Baumberg. Alcohol in Europe. Institute of Alcohol Studies for the European Commission, London. Public Health Perspective: Report summary, p. 6.
<http://dse.univr.it/addiction/documents/External/alcoholineu.pdf>. *Drugs Education Prev Policy* 13, 483-488, 2006.
- Anderson P, A Cremona, A Paton, C Turner, P Wallace: The risk of alcohol. *Addiction* 1993; 88(11):1493-1508.
- Andreu V, A Mas, M Bruguera, JM Salmeron, V Moreno, S Nogue, J Rodes: Ecstasy: a common cause of severe acute hepatotoxicity. *J Hepatol* 1998; 29(3):394-397.
- Anonymous: Carcinoma of the prostate. *N Engl J Med* 1991; 324:1892-1893.
- Arditti J. Ketamine, déviation d'usage en France, CEIP, Marseille, France. 2000.

- Arnsten JH, PA Demas, H Farzadegan, RW Grant, MN Gourevitch, CJ Chang, D Buono, H Eckholdt, AA Howard, EE Schoenbaum: Antiretroviral therapy adherence and viral suppression in HIV-infected drug users: comparison of self-report and electronic monitoring. *Clin Infect Dis* 2001; 33(8):1417-1423.
- Ashton CH: Pharmacology and effects of cannabis: a brief review. *Br J Psychiatry* 2001; 178:101-106.
- Baan R, K Straif, Y Grosse, B Secretan, F El Ghissassi, V Bouvard, A Altieri, V Coglianò. Carcinogenicity of alcoholic beverages. WHO International Agency for Research on Cancer Monograph Working Group. *Lancet Oncol* 8, 292-293, 2007.
- Bagasra O, RJ Pomerantz: Human immunodeficiency virus type 1 replication in peripheral blood mononuclear cells in the presence of cocaine. *J Infect Dis* 1993; 168(5):1157-1164.
- Bagatell CJ, WJ Bremner: Androgens in men--uses and abuses. *N Engl J Med* 1996; 334(11):707-714.
- Bagia S, PM Hewitt, DL Morris: Anabolic steroid-induced hepatic adenomas with spontaneous haemorrhage in a bodybuilder. *Aust N Z J Surg* 2000; 70(9):686-687.
- Balakrishnan R, S Allender, P Scarborough, P Webster, M Rayner: The burden of alcohol-related ill health in the United Kingdom. *J Public Health (Oxf)* 2009; 31(3):366-373.
- Baldwin GC, MD Roth, DP Tashkin: Acute and chronic effects of cocaine on the immune system and the possible link to AIDS. *J Neuroimmunol* 1998; 83(1-2):133-138.
- Ball JC, WR Lange, CP Myers, SR Friedman: Reducing the risk of AIDS through methadone maintenance treatment. *J Health Soc Behav* 1988; 29(3):214-226.
- Barendrecht JJ, L Bonneux, PJ van der Maas. The health care costs of smoking. *N Engl J Med* 337, 1052-1057, 1997.
- Bargagli AM, M Hickman, M Davoli, CA Perucci, P Schifano, M Buster, T Brugal, J Vicente: Drug-related mortality and its impact on adult mortality in eight European countries. *Eur J Public Health* 2006; 16(2):198-202.
- Barsky SH, MD Roth, EC Kleerup, M Simmons, DP Tashkin: Histopathologic and molecular alterations in bronchial epithelium in habitual smokers of marijuana, cocaine, and/or tobacco. *J Natl Cancer Inst* 1998; 90(16):1198-1205.
- Bashour TT: Acute myocardial infarction resulting from amphetamine abuse: a spasm-thrombus interplay? *Am Heart J* 1994; 128(6 Pt 1):1237-1239.
- Bauman JL, JJ Grawe, AP Winecoff, RJ Hariman: Cocaine-related sudden cardiac death: a hypothesis correlating basic science and clinical observations. *J Clin Pharmacol* 1994; 34(9):902-911.
- Becker U, M Gronbaek, D Johansen, ThIA Sorensen: Lower risk for alcohol-induced cirrhosis in wine drinkers. *Hepatology* 2002; 35(4):868-875.
- Beebe DK, E Walley: Smokable methamphetamine ('ice'): an old drug in a different form. *Am Fam Physician* 1995; 51(2):449-453.
- Block RI, R Farinpour, JA Schlechte: Effects of chronic marijuana use on testosterone, luteinizing hormone, follicle stimulating hormone, prolactin and cortisol in men and women. *Drug Alcohol Depend* 1991; 28:121-128.
- Bloom JW, WT Kaltenborn, P Paoletti, A Camilli, MD Lebowitz: Respiratory effects of non-tobacco cigarettes. *Br Med J (Clin Res Ed)* 1987; 295(6612):1516-1518.

- Bloor M, M Gannon, G Hay, G Jackson, AH Leyland, N McKeganey: Contribution of problem drug users' deaths to excess mortality in Scotland: secondary analysis of cohort study. *BMJ* 2008; 337:a478.
- Blot WJ, JK McLaughlin, DM Winn, DF Austin, RF Greenberg, S Preston-Martin, L Bernstein, JB Schoenberg, A Stemhagen, JF Fraumeni. Smoking and drinking in relation to oral and pharyngeal cancer. *Cancer Res* 48, 3282-3287, 1988.
- Boothby LA, PL Doering: Buprenorphine for the treatment of opioid dependence. *Am J Health Syst Pharm* 2007; 64(3):266-272.
- Bostwick MJ, TW Lineberry. The 'meth' epidemic: managing acute psychosis, agitation, and suicide risk. *J Fam Prac* 5, 46-62, 2006.
- Boyadjiev NP, KN Georgieva, RI Massaldjieva, SI Gueorguiev: Reversible hypogonadism and azoospermia as a result of anabolic-androgenic steroid use in a bodybuilder with personality disorder. A case report. *J Sports Med Phys Fitness* 2000; 40(3):271-274.
- Brands B, B Sproule, J Marshman. Cocaine. In B. Brands, B. Sproule & J. Marshman (Eds.), *Drug & Drug Abuse* (3rd ed., pp. 239-248). Toronto: Addiction Research Foundation. 1988.
- Brauer RB, CD Heidecke, W Nathrath, KT Beckurts, P Vorwald, TR Zilker, U Schweigart, AH Holscher, JR Siewert: Liver transplantation for the treatment of fulminant hepatic failure induced by the ingestion of ecstasy. *Transpl Int* 1997; 10(3):229-233.
- Brody SL, CM Slovis, KD Wrenn: Cocaine-related medical problems: consecutive series of 233 patients. *Am J Med* 1990; 88(4):325-331.
- Brower KJ: Anabolic steroid abuse and dependence. *Curr Psychiatry Rep* 2002; 4(5):377-387.
- Brust JC, RW Richter: Stroke associated with addiction to heroin. *J Neurol Neurosurg Psychiatry* 1976; 39(2):194-199.
- Buchanan JF, CR Brown: 'Designer drugs'. A problem in clinical toxicology. *Med Toxicol Adverse Drug Exp* 1988; 3(1):1-17.
- Buck ML, JL Blumer: Opioids and other analgesics. Adverse effects in the intensive care unit. *Crit Care Clin* 1991; 7(3):615-637.
- Bullen C: Impact of tobacco smoking and smoking cessation on cardiovascular risk and disease. *Expert Rev Cardiovasc Ther* 2008; 6(6):883-895.
- Busch FW, DA Seid, ET Wei: Mutagenic activity of marijuana smoke condensates. *Cancer Lett* 1979; 6(6):319-324.
- Buster M, L Rook, GHA van Brussel, J van Ree, W van den Brink: Chasing the dragon, related to the impaired lung function among heroin users. *Drug Alcohol Depend* 2002; 68(2):221-228.
- Buxton N, NS McConachie: Amphetamine abuse and intracranial haemorrhage. *J R Soc Med* 2000; 93(9):472-477.
- Calignano A, I Katona, F Desarnaud, A Giuffrida, G La Rana, K Mackie, TF Freund, D Piomelli: Bidirectional control of airway responsiveness by endogenous cannabinoids. *Nature* 2000; 408(6808):96-101.
- Callaway CW, RF Clark: Hyperthermia in psychostimulant overdose. *Ann Emerg Med* 1994; 24(1):68-76.
- Campsmith ML, AK Nakashima, JL Jones: Association between crack cocaine use and high-risk sexual behaviors after HIV diagnosis. *J Acquir Immune Defic Syndr* 2000; 25(2):192-198.
- Caplan GA, BA Brigham: Marijuana smoking and carcinoma of the tongue. Is there an association? *Cancer* 1990; 66(5):1005-1006.
- Caplehorn JR, MS Dalton, F Haldar, AM Petrenas, JG Nisbet: Methadone maintenance and addicts' risk of fatal heroin overdose. *Subst Use Misuse* 1996b; 31(2):177-196.

- Caplehorn JR, MS Dalton, F Haldar, AM Petrenas, JG Nisbet: Methadone maintenance and addicts' risk of fatal heroin overdose. *Subst Use Misuse* 1996a; 31(2):177-196.
- Carter EF: Dental implications of narcotic addiction. *Aust Dent J* 1978; 23(4):308-310.
- Chan PC, RC Sills, AG Braun, JK Haseman, JR Bucher: Toxicity and carcinogenicity of delta 9-tetrahydrocannabinol in Fischer rats and B6C3F1 mice. *Fundam Appl Toxicol* 1996; 30(1):109-117.
- Cheng TO: Risk of acute myocardial infarction in cocaine abusers. *Circulation* 2000; 101(23):E227.
- Chernoff RW, MH Wallen, OF Muller: Cardiac toxicity of methylphenidate. Report of two cases. *N Engl J Med* 1962; 266:400-401.
- Chin MY, RA Kreutzer, JE Dyer. Acute poisoning from gamma-hydroxybutyrate in California. *West J Med* 156, 380-384, 1992.
- Chin RL, KA Sporer, B Cullison, JE Dyer, TD Wu. Clinical course of gamma-hydroxybutyrate overdose. *Ann Emerg Med* 31, 716-722, 1998.
- Cho CM, R Hirsch, Johnstone S: General and oral health implications of cannabis use. *Aust Dent J* 2005; 50(2):70-74.
- Cho E, SA Smith-Warner, J Ritz, PA van den Brandt, GA Colditz, AR Folsom, JL Freudenheim, E Giovannucci, RA Goldbohm, S Graham, L Holmberg, DH Kim, N Malila, AB Miller, P Pietinen, ThE Rohan, ThA Sellers, FE Speizer, WC Willett, A Wolk, DJ Hunter: Alcohol intake and colorectal cancer: a pooled analysis of 8 cohort studies. *Ann Intern Med* 2004; 140(8):603-613.
- Chu PS-K, WK Ma, SC-W Wong, RW-H Chu, CH Cheng, S Wong, JM-L Tse, FL Lau, MK Yiu, CW Man: The destruction of the lower urinary tract by ketamine abuse: a new syndrome? *BJU Int* 2008; 102(11):1616-1622.
- Chugh SS, C Socoteanu, K Reinier, J Waltz, J Jui, K Gunson: A community-based evaluation of sudden death associated with therapeutic levels of methadone. *Am J Med* 2008; 121(1):66-71.
- Cicero TJ, RD Bell, WG Wiest, JH Allison, K Polakoski, E Robins: Function of the male sex organs in heroin and methadone users. *N Engl J Med* 1975; 292(17):882-887.
- Citron BP, M Halpern M, M McCarron, GD Lundberg, R McCormick, IJ Pincus, D Tatter, BJ Haverback: Necrotizing angitis associated with drug abuse. *N Engl J Med* 1970; 283(19):1003-1011.
- Coffin PO, S Galea, J Ahern, AC Leon, D Vlahov, K Tardiff: Opiates, cocaine and alcohol combinations in accidental drug overdose deaths in New York City, 1990-98. *Addiction* 2003; 98(6):739-747.
- Cohen EJ, AL Klatsky, MA Armstrong: Alcohol use and supraventricular arrhythmia. *Am J Cardiol* 1988; 62(13):971-973.
- Cohen MM, Shiloh Y: Genetic toxicology of lysergic acid diethylamide (LSD-25). *Mutat Res* 1977; 47(3-4):183-209.
- Colebunders B, P van Erps: Cystitis due to the use of ketamine as a recreational drug: a case report. *J Med Case Reports* 2008; 2:219.
- Connock M, A Juarez-Garcia, S Jowett, E Frew, Z Liu, RJ Taylor, A Fry-Smith, E Day, N Lintzeris, T Roberts, A Burls, RS Taylor: Methadone and buprenorphine for the management of opioid dependence: a systematic review and economic evaluation. *Health Technol Assess* 2007; 11:i-iv.
- Cook JA, JK Burke-Miller, MH Cohen, RL Cook, D Vlahov, TE Wilson, ET Golub, RM Schwartz, AA Howard, C Ponath, MW Plankey, AM Levine, DD Grey: Crack cocaine, disease progression, and mortality in a multicenter cohort of HIV-1 positive women. *AIDS* 2008; 22(11):1355-1363.
- Cornish JW, CP O'Brien: Crack cocaine abuse: an epidemic with many public health consequences. *Annu Rev Public Health* 1996; 17:259-273.

- Corrao G, V Bagnardi, A Zambon, S Arico: Exploring the dose-response relationship between alcohol consumption and the risk of several alcohol-related conditions: a meta-analysis. *Addiction* 1999; 94(10):1551-1573.
- Corrao G, V Bagnardi, A Zambon, C La Vecchia: A metaanalysis of alcohol consumption and the risk of 15 diseases. *Prev Med* 38, 613-619, 2004.
- Cottrell AM, D Gillatt: Consider ketamine misuse in patients with urinary symptoms. *Practitioner* 2008; 252(1711):5.
- Council for Public Health and Care and Council for Social Development [Raad voor de Volksgezondheid en Zorg and Raad voor de Maatschappelijke Ontwikkeling]. *Verslavingszorg herijkt*, juli 1999. 1999.
- Creagh TM, A Rubin, DJ Evans: Hepatic tumours induced by anabolic steroids in an athlete. *J Clin Pathol* 1988; 41(4):441-443.
- Crean RD, NA Crane, BJ Mason: An evidence based review of acute and long-term effects of cannabis use on executive cognitive functions. *J Addict Med* 2011; 5(1):1-8.
- Cruts AAN, M Buster, J Vicente, I Deerenberg, M van Laar: Estimating the Total Mortality Among Problem Drug Users. *Subst Use Misuse* 43, 733-747, 2008.
- Cunningham CO, NL Sohler, KM Berg, S Shapiro, D Heller: Type of substance use and access to HIV-related health care. *AIDS Patient Care STDS* 2006; 20(6):399-407.
- Cunningham EE, MA Zielezny, RC Venuto: Heroin-associated nephropathy. A nationwide problem. *JAMA* 1983; 250(21):2935-2936.
- Curtis EK: Meth mouth: a review of methamphetamine abuse and its oral manifestations. *Gen Dent* 2006; 54(2):125-129.
- Cygan J, M Trunsky, T Corbridge: Inhaled heroin-induced status asthmaticus: five cases and a review of the literature. *Chest* 2000; 117(1):272-275.
- D'Andrea A, P Caso, G Salerno, R Scarafile, G De Corato, C Mita, G Di Salvo, S Severino, S Cuomo, B Liccardo, N Esposito, R Calabro: Left ventricular early myocardial dysfunction after chronic misuse of anabolic androgenic steroids: a Doppler myocardial and strain imaging analysis. *Br J Sports Med* 2007; 41(3):149-155.
- Daccarett M, M Freih, C Machado: Acute cannabis intoxication mimicking brugada-like ST segment abnormalities. *Int J Cardiol* 2007; 119:235-236.
- Dackis CA, M Gurpegui, AL Pottash, MS Gold: Methadone induced hypoadrenalism. *Lancet* 1982; 2(8308):1167.
- Dalgarno PJ, D Shewan: Illicit use of ketamine in Scotland. *J Psychoactive Drugs* 1996; 28(2):191-199.
- Darke S, S Kaye, R McKetin, J Duflou. Physical and psychological harms of psychostimulant use. TR-286. NDARC. 2007a.
- Darke S, J Ross, A Williamson, KL Mills, A Havard, M Teesson: Patterns and correlates of attempted suicide by heroin users over a 3-year period: findings from the Australian treatment outcome study. *Drug Alcohol Depend* 2007b; 87(2-3):146-152.
- Darke S, M Torok, S Kaye, J Ross: Attempted suicide, self-harm, and violent victimization among regular illicit drug users. *Suicide Life Threat Behav* 2010; 40(6):587-596.
- Davis RM, M Wakefield, A Amos, PC Gupta: The Hitchhiker's Guide to Tobacco Control: a global assessment of harms, remedies, and controversies. *Annu Rev Public Health* 2007; 28:171-194.
- De la Torre R, M Farre, PN Roset, N Pizarro, S Abanades, M Segura, J Segura, J Cami: Human pharmacology of MDMA: pharmacokinetics, metabolism, and disposition. *Ther Drug Monit* 2004c; 26(2):137-144.

- De la Torre R, M Farre, PN Roset, N Pizarro, S Abanades, M Segura, J Segura, J Cami: Human pharmacology of MDMA: pharmacokinetics, metabolism, and disposition. *Ther Drug Monit* 2004; 26(2):137-144.
- Degenhardt L, W Hall, M Warner-Smith, M Lynskey. Illicit drug use. M Ezzati, A Lopez, and C Murray. Comparative risk assessment. [1], 1109-1176. 2004. Geneva, WHO (World Health Organisation). The World Health Report 2003: Shaping the Future. 2004.
- Delaney K, RS Hoffman: Pulmonary infarction associated with crack cocaine use in a previously healthy 23-year-old woman. *Am J Med* 1991; 91:92-94.
- Den Dekker J. Behandelpanning in de tandartspraktijk. Thesis. University of Amsterdam 1990. 1990.
- Dettmeyer R, P Schmidt, F Musshoff, C Dreisvogt, B Madea: Pulmonary edema in fatal heroin overdose: immunohistological investigations with IgE, collagen IV and laminin - no increase of defects of alveolar-capillary membranes. *Forensic Sci Int* 2000; 110(2):87-96.
- Dewit DJ, J Hance, DR Offord, A Ogborne: The influence of early and frequent use of marijuana on the risk of desistance and of progression to marijuana-related harm. *Prev Med* 2000; 31(5):455-464.
- Dimba EAO, BT Gjertsen, T Bredholt, KO Fossan, DE Costea, GW Francis, AC Johannessen, OK Vintermyr: Khat (*Catha edulis*)-induced apoptosis is inhibited by antagonists of caspase-1 and -8 in human leukaemia cells. *Br J Cancer* 2004; 91(9):1726-1734.
- Dishotsky NI, WD Loughman, RE Mogar, WR Lipscomb: LSD and genetic damage. *Science* 1971; 172(982):431-440.
- Doll R: Epidemiological evidence of the effects of behaviour and the environment on the risk of human cancer. *Recent Results Cancer Res* 1998; 154:3-21.
- Drummer OH, DL Ranson: Sudden death and benzodiazepines. *Am J Forensic Med Pathol* 1996; 17(4):336-342.
- Drummer OH, M Syrjanen, K Opeskin, S Cordner: Deaths of heroin addicts starting on a methadone maintenance programme. *Lancet* 1990; 335(8681):108.
- Dubrow A, N Mittman, V Ghali, W Flamenbaum: The changing spectrum of heroin-associated nephropathy. *Am J Kidney Dis* 1985; 5(1):36-41.
- Eagle KA, EM Isselbacher, RW DeSanctis: Cocaine-related aortic dissection in perspective. *Circulation* 2002; 105(13):1529-1530.
- Ebadi M, RF Pfeiffer, LC Murrin: Pathogenesis and treatment of neuroleptic malignant syndrome. *Gen Pharmacol* 1990; 21(4):367-386.
- Edston E, M Hage-Hamsten: Anaphylactoid shock--a common cause of death in heroin addicts? *Allergy* 1997; 52(9):950-954.
- El Shoura SM, M Abdel Aziz M, ME Ali, MM el Said, KZ Ali, MA Kemei, AM Raoof, M Allam, EM Elmalik: Deleterious effects of khat addiction on semen parameters and sperm ultrastructure. *Hum Reprod* 1995; 10(9):2295-2300.
- El-Omar MM, K Ray, R Geary: Intracerebral haemorrhage in a young adult: consider amphetamine abuse. *Br J Clin Pract* 1996; 50(2):115-116.
- Ellis AJ, JA Wendon, B Portmann, R Williams: Acute liver damage and ecstasy ingestion. *Gut* 1996; 38(3):454-458.
- Ellison RC, Y Zhang, CE McLennan, KJ Rothman: Exploring the relation of alcohol consumption to risk of breast cancer. *Am J Epidemiol* 2001; 154(8):740-747.
- EMCDDA: European Monitoring Centre for Drugs and Drug Addiction (EMCDDA). Annual report 2010. The state of the drugs problem in Europe. Luxembourg, Office for Official Publications of the European Communities. 2010.

- Ettinger NA, RJ Albin: A review of the respiratory effects of smoking cocaine. *Am J Med* 1989; 87(6):664-668.
- Ettinger PO, CF Wu, CJ De La Cruz, AB Weisse, SS Ahmed, TJ Regan: Arrhythmias and the 'Holiday Heart': alcohol-associated cardiac rhythm disorders. *Am Heart J* 1978; 95(5):555-562.
- EUPHIX: Medical conditions and social harms related to alcohol consumption. In: EUPHIX, EUphact. RIVM, Bilthoven, The Netherlands. http://www.euphix.org/object_class/euph_health_behaviours_alcohol_use.html. 2007.
- Evans NA: Current concepts in anabolic-androgenic steroids. *Am J Sports Med* 2004; 32(2):534-542.
- Extein I: Methylphenidate-induced choreoathetosis. *Am J Psychiatry* 1978; 135(2):252-253.
- Ferenchick G, D Schwartz, M Ball, K Schwartz: Androgenic-anabolic steroid abuse and platelet aggregation: a pilot study in weight lifters. *Am J Med Sci* 1992; 303(2):78-82.
- Fidler H, A Dhillon, D Gertner, A Burroughs: Chronic ecstasy (3,4-methylenedioxymetamphetamine) abuse: a recurrent and unpredictable cause of severe acute hepatitis. *J Hepatol* 1996; 25(4):563-566.
- Filmore KM, WC Kerr, T Stockwell, T Chikritzhs, A Bostrom: Moderate alcohol use and reduced mortality risk: systematic error in prospective studies. *Addict Res Theory* 14, 101-132, 2006.
- Fineschi V, I Riezzo, F Centini, E Silingardi, M Licata, G Beduschi, SB Karch: Sudden cardiac death during anabolic steroid abuse: morphologic and toxicologic findings in two fatal cases of bodybuilders. *Int J Legal Med* 2007; 121(1):48-53.
- Fischer B, MF Cruz, J Rehm: Illicit opioid use and its key characteristics: a select overview and evidence from a Canadian multisite cohort of illicit opioid users (OPICAN). *Can J Psychiatry* 2006; 51(10):624-634.
- Fischer B, E Haydon, J Rehm, M Kraijden, J Reimer: Injection drug use and the hepatitis C virus: considerations for a targeted treatment approach--the case study of Canada. *J Urban Health* 2004; 81(3):428-447.
- Fossey MD, MB Hamner: Clonazepam-related sexual dysfunction in male veterans with PTSD. *Anxiety* 1994; 1(5):233-236.
- Frankle MA, R Eichberg, SB Zachariah: Anabolic androgenic steroids and a stroke in an athlete: case report. *Arch Phys Med Rehabil* 1988b; 69(8):632-633.
- Frankle MA, R Eichberg, SB Zachariah: Anabolic androgenic steroids and a stroke in an athlete: case report. *Arch Phys Med Rehabil* 1988a; 69(8):632-633.
- Fredericks RK, DS Lefkowitz, VR Challa, BT Troost: Cerebral vasculitis associated with cocaine abuse. *Stroke* 1991; 22(11):1437-1439.
- Friedlander AH, DA Gorelick: Dental management of the cocaine addict. *Oral Surg Oral Med Oral Pathol* 1988; 65(1):45-48.
- Friedman SA, SE Hirsch: Extreme hyperthermia after LSD ingestion. *JAMA* 1971; 217(11):1549-1550.
- Fuchs CS, MJ Stampfer, GA Colditz, EL Giovannucci, JE Manson, I Kawachi, DJ Hunter, SE Hankinson, CH Hennekens, B Rosner: Alcohol consumption and mortality among women. *N Engl J Med* 1995; 332(19):1245-1250.
- Garbino J, JA Henry, G Mentha, JA Romand: Ecstasy ingestion and fulminant hepatic failure: liver transplantation to be considered as a last therapeutic option. *Vet Hum Toxicol* 2001; 43(2):99-102.
- Gezondheidsraad: Volksgezondheidsschade door passief roken. 2003/21. 2008.

- Ghadirian AM, L Annable, MC Belanger: Lithium, benzodiazepines, and sexual function in bipolar patients. *Am J Psychiatry* 1992; 149(6):801-805.
- Gibson A, D Randall, L Degenhardt: The increasing mortality burden of liver disease among opioid dependent people: Cohort study. *Addiction* 2011.
- Goldenberg D, J Lee, WM Koch, MM Kim, B Trink, D Sidransky, CS Moon: Habitual risk factors for head and neck cancer. *Otolaryngol Head Neck Surg* 2004; 131(6):986-993.
- Goppel MA, W van den Brink, JM van Ree : ['Pharmacologic interventions in drug addiction': an advisory report from the Health Council of the Netherlands]. 'Medicamenteuze interventies bij drugverslaving'; een advies van de Gezondheidsraad. *Ned Tijdschr Geneeskd* 2003; 147(34):1637-1640.
- Gorayski P, CH Thompson, HS Subhash, AC Thomas: Hepatocellular carcinoma associated with recreational anabolic steroid use. *Br J Sports Med* 2008; 42(1):74-75.
- Gordon CJ, WP Watkinson, JP O'Callaghan, DB Miller: Effects of 3,4-methylenedioxymethamphetamine on autonomic thermoregulatory responses of the rat. *Pharmacol Biochem Behav* 1991; 38(2):339-344.
- Gore SM: Fatal uncertainty: death-rate from use of ecstasy or heroin. *Lancet* 1999; 354(9186):1265-1266.
- Goudie AJ: Comparative effects of cathinone and amphetamine on fixed-interval operant responding: a rate-dependency analysis. *Pharmacol Biochem Behav* 1985; 23(3):355-365.
- Gouzoulis-Mayfrank E, J Daumann: Neurotoxicity of methylenedioxyamphetamines (MDMA; ecstasy) in humans: how strong is the evidence for persistent brain damage? *Addiction* 2006; 101(3):348-361.
- Gouzoulis-Mayfrank E, B Thelen, E Habermeyer, HJ Kunert, KA Kovar, H Lindenblatt, L Hermlle, M Spitzer, H Sass: Psychopathological, neuroendocrine and autonomic effects of 3,4-methylenedioxyethylamphetamine (MDE), psilocybin and d-methamphetamine in healthy volunteers. Results of an experimental double-blind placebo-controlled study. *Psychopharmacology (Berl)* 1999; 142(1):41-50.
- Greenhill LL, J Puig-Antich, H Novacenko, M Solomon, C Anghern, J Florea, R Goetz, B Fiscina, EJ Sachar: Prolactin, growth hormone and growth responses in boys with attention deficit disorder and hyperactivity treated with methylphenidate. *J Am Acad Child Psychiatry* 1984; 23(1):58-67.
- Greenhill LL, RL Findling, JM Swanson: A double-blind, placebo-controlled study of modified-release methylphenidate in children with attention-deficit/hyperactivity disorder. *Pediatrics* 2002; 109(3):E39.
- Grobbee DE, EB Rim, U Keil, S Renaud: Alcohol and the Cardiovascular System. In: Macdonald I, editor. *Health issues related to alcohol consumption*. Cornwall: Blackwell Science Ltd. 125-179 (1999). 1999.
- Grotenhermen F: The toxicology of cannabis and cannabis prohibition. *Chem Biodivers* 2007; 4(8):1744-1769.
- Guzman M: Cannabinoids: potential anticancer agents. *Nat Rev Cancer* 2003; 3(10):745-755.
- Haemmig RB, W Tschacher: Effects of high-dose heroin versus morphine in intravenous drug users: a randomised double-blind crossover study. *J Psychoactive Drugs* 2001; 33(2):105-110.
- Haim DY, ML Lippmann, SK Goldberg, MD Walkenstein: The pulmonary complications of crack cocaine. A comprehensive review. *Chest* 1995a; 107(1):233-240.

- Hakim LY: Influence of khat on seminal fluid among presumed infertile couples. *East Afr Med J* 2002; 79(1):22-28.
- Halbach H: Medical aspects of the chewing of khat leaves. *Bull World Health Organ* 1972; 47(1):21-29.
- Hall AP, JA Henry: Acute toxic effects of 'Ecstasy' (MDMA) and related compounds: overview of pathophysiology and clinical management. *Br J Anaesth* 2006a; 96(6):678-685.
- Hall W: The respiratory risks of cannabis smoking. *Addiction* 1998; 93(10):1461-1463.
- Hall W, C Doran, L Degenhardt, D Shephard: Illicit Opiate Abuse. In: DT Jamison, JG Breman, AR Measham, et al., eds. *Disease Control Priorities in Developing Countries*. Washington (DC): Worldbank, 2006. 2006b.
- Hall , N Solowij: Adverse effects of cannabis. *Lancet* 1998; 352(9140):1611-1616.
- Hall W, N Solowij, J Lemon: The health and psychological consequences of cannabis use. *National Drug Strategy Monograph Series*, 25. <http://www.health.gov.au/internet/main/publishing.nsf/Content/health-pubs-drug-cannab2-ch46.htm>. 1994.
- Hall W, M Christie, D Currow: Cannabinoids and cancer: causation, remediation, and palliation. *Lancet Oncol* 2005; 6(1):35-42.
- Hall W, L Degenhardt: Adverse health effects of non-medical cannabis use. *Lancet* 2009; 374(9698):1383-1391.
- Hamajima N, K Hirose, K Tajima, T Rohan, EE Calle, CW Heath, Jr., RJ Coates, JM Liff, R Talamini: Alcohol, tobacco and breast cancer--collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with breast cancer and 95,067 women without the disease. *Br J Cancer* 2002; 87(11):1234-1245.
- Hamamoto DT, NL Rhodus: Methamphetamine abuse and dentistry. *Oral Dis* 2009; 15(1):27-37.
- Hamilton D: Heat stroke. *Anaesthesia* 32, 271, 1976.
- Harari MD, D Netzer: Genital examination under ketamine sedation in cases of suspected sexual abuse. *Arch Dis Child* 1994; 70(3):197-198.
- Hartgens F, H Kuipers: Effects of androgenic-anabolic steroids in athletes. *Sports Med* 2004; 34(8):513-554.
- Harzke AJ, ML Williams, AM Bowen: Binge use of crack cocaine and sexual risk behaviors among African-American, HIV-positive users. *AIDS Behav* 2009; 13(6):1106-1118.
- Hashibe M, H Morgenstern, Y Cui, DP Tashkin, ZF Zhang, W Cozen, TM Mack, S Greenland: Marijuana use and the risk of lung and upper aerodigestive tract cancers: results of a population-based case-control study. *Cancer Epidemiol Biomarkers Prev* 2006; 15(10):1829-1834.
- Hashibe M, K Straif, DP Tashkin, H Morgenstern, S Greenland, ZF Zhang: Epidemiologic review of marijuana use and cancer risk. *Alcohol* 2005; 35(3):265-275.
- Hasler F, U Grimberg, MA Benz, T Huber, FX Vollenweider: Acute psychological and physiological effects of psilocybin in healthy humans: a double-blind, placebo-controlled dose-effect study. *Psychopharmacology (Berl)* 2004; 172(2):145-156.
- Hassan NA, AA Gunaid, AA bdo-Rabbo, ZY bdel-Kader, MA al-Mansoob, AY Awad, IM Murray-Lyon: The effect of Qat chewing on blood pressure and heart rate in healthy volunteers. *Trop Doct* 2000; 30(2):107-108.
- Hatsukami DK, LF Stead, PC Gupta: Tobacco addiction. *Lancet* 371, 2027-2038, 2008.

- Health Council of the Netherlands: Ethanol (ethyl alcohol); Evaluation of the health effects from occupational exposure. The Hague: Health Council of the Netherlands, publication no. 2006/06OSH. 2006.
- Hechtman L, B Greenfield: Long-term use of stimulants in children with attention deficit hyperactivity disorder: safety, efficacy, and long-term outcome. *Paediatr Drugs* 2003; 5(12):787-794.
- Heesch CM, CR Wilhelm, J Ristich, J Adnane, FA Bontempo, WR Wagner: Cocaine activates platelets and increases the formation of circulating platelet containing microaggregates in humans. *Heart* 2000; 83(6):688-695.
- Heil SH, GJ Badger, ST Higgins: Alcohol dependence among cocaine-dependent outpatients: demographics, drug use, treatment outcome and other characteristics. *J Stud Alcohol* 2001; 62(1):14-22.
- Hendrickson RG, R Cloutier, KJ McConnell: Methamphetamine-related emergency department utilization and cost. *Acad Emerg Med* 2008; 15(1):23-31.
- Hendriks VM, W van den Brink, JM van Ree, P Blanken, M Koeter, BJ van Zwieten: Heroïne en methadon op medisch voorschrift voor chronisch, therapieresistente heroïneverslaafden: twee gerandomiseerde studies. *Ned Tijdschr Geneesk* 147, 2065-2071, 2003.
- Henning RJ, LD Wilson, JM Glauser: Cocaine plus ethanol is more cardiotoxic than cocaine or ethanol alone. *Crit Care Med* 1994; 22(12):1896-1906.
- Henry JA: Ecstasy and the dance of death. *BMJ* 1992; 305(6844):5-6.
- Her M, J Rehm: Alcohol and all-cause mortality in Europe 1982-1990: a pooled cross-section time-series analysis. 93, 1335-1340, 1998. *Addiction* 93, 1335-1340, 1998.
- Hickman M, Z Carnwath, P Madden, M Farrell, C Rooney, R Ashcroft, A Judd, G Stimson: Drug-related mortality and fatal overdose risk: pilot cohort study of heroin users recruited from specialist drug treatment sites in London. *J Urban Health* 2003; 80(2):274-287.
- Hill CM, A Gibson: The oral and dental effects of q'at chewing. *Oral Surg Oral Med Oral Pathol* 1987; 63(4):433-436.
- Hinkin CH, TR Barclay, SA Castellon, AJ Levine, RS Durvasula, SD Marion, HF Myers, D Longshore: Drug use and medication adherence among HIV-1 infected individuals. *AIDS Behav* 2007; 11(2):185-194.
- Ho EL, SA Josephson, HS Lee, WS Smith: Cerebrovascular complications of methamphetamine abuse. *Neurocrit Care* 2009; 10(3):295-305.
- Hoffmann D, KD Brunnemann, GB Gori, EL Wynder: On the carcinogenicity of marijuana smoke. *Runeckles, V.C. ed. 63-81. Recent Advances in Phytochemistry. 1975.*
- Hollander JE, RS Hoffman, JL Burstein, RD Shih, HCJ Thode: Cocaine-associated myocardial infarction. Mortality and complications. Cocaine-Associated Myocardial Infarction Study Group. *Arch Intern Med* 1995; 155(10):1081-1086.
- Hollister LE: Health aspects of cannabis. *Pharmacol Rev* 1986a; 38(1):1-20.
- Hollister LE: Marijuana and immunity. *J Psychoactive Drugs* 1992; 24(2):159-164.
- Holman CD, DR English, E Milne, MG Winter: Meta-analysis of alcohol and all-cause mortality: a validation of NHMRC recommendations. *Med J Aust* 1996; 164(3):141-145.
- Holtkamp K, B Peters-Wallraf, S Wuller, R Pfaaffle, B Herpertz-Dahlmann: Methylphenidate-related growth impairment. *J Child Adolesc Psychopharmacol* 2002; 12(1):55-61.
- Hsairi M, N Achour, B Zouari, H Ben Romdhane, A Achour, M Maalej, T Nacef: [Etiologic factors in primary bronchial carcinoma in Tunisia]

- Facteurs etiologiques du cancer bronchique primitif en Tunisie. *Tunis Med* 1993; 71(5):265-268.
- Hughes S, PM Calverley: Heroin inhalation and asthma. *BMJ* 1988; 297(6662):1511-1512.
- Hurley BF, DR Seals, JM Hagberg, AC Goldberg, SM Ostrove, JO Holloszy, WG Wiest, AP Goldberg: High-density-lipoprotein cholesterol in bodybuilders v powerlifters. Negative effects of androgen use. *JAMA* 1984; 252:507-513.
- Hylkema MN, PJ Sterk, WI de Boer, DS Postma: Tobacco use in relation to COPD and asthma. *Eur Respir J* 2007; 29(3):438-445.
- Ibrahim MM, H Deng, A Zvonok, DA Cockayne, J Kwan, HP Mata, TW Vanderah, J Lai, F Porreca, A Makriyannis, TPJ Malan: Activation of CB2 cannabinoid receptors by AM1241 inhibits experimental neuropathic pain: pain inhibition by receptors not present in the CNS. *Proc Natl Acad Sci U S A* 2003; 100(18):10529-10533.
- ICAP: International Center for alcohol policies (ICAP). Reports 17. The structure of the beverage alcohol industry. <http://www.icap.org/Publications/ICAPReports/tabid/75/Default.aspx>. 2006.
- INSERM: Expertises collectives: Tabac - Comprendre la dépendance pour agir, pp 1-473. 2004.
- Ishak KG, HJ Zimmerman: Hepatotoxic effects of the anabolic/androgenic steroids. *Semin Liver Dis* 7, 230-236, 1987.
- Islam MN, HKuroki, B Hongcheng, Y Ogura, N Kawaguchi, S Onishi, C Wakasugi: Cardiac lesions and their reversibility after long term administration of methamphetamine. *Forensic Sci Int* 1995; 75(1):29-43.
- Islam MW, OA al Shabanah, MM al Harbi, NM al Gharably: Evaluation of teratogenic potential of khat (*Catha edulis* Forsk.) in rats. *Drug Chem Toxicol* 1994; 17(1):51-68.
- Israel JA, K Lee: Amphetamine usage and genital self-mutilation. *Addiction* 2002; 97(9):1215-1218.
- Iversen L: Speed, ecstasy and ritalin: the science of amphetamines. Oxford: Oxford University Press. 2008.
- Jacob V, S Vellappally, J Smejkalova: The influence of cigarette smoking on various aspects of periodontal health. *Acta Medica (Hradec Kralove)* 2007; 50(1):3-5.
- Jaehne EJ, A Salem, RJ Irvine: Effects of 3,4-methylenedioxymethamphetamine and related amphetamines on autonomic and behavioral thermoregulation. *Pharmacol Biochem Behav* 2005; 81(3):485-496.
- Jaffe JA, PL Kimmel: Chronic nephropathies of cocaine and heroin abuse: a critical review. *Clin J Am Soc Nephrol* 2006; 1(4):655-667.
- Jaffe RB: Cardiac and vascular involvement in drug abuse. *Semin Roentgenol* 1983; 18(3):207-212.
- Jamal M, Z Saadi, TR Morgan: Alcohol and hepatitis C. *Dig Dis* 2005; 23(3-4):285-296.
- Jansen KL: A review of the nonmedical use of ketamine: use, users and consequences. *J Psychoactive Drugs* 2000; 32(4):419-433.
- Jin B, L Turner, WA Walters, DJ Handelsman: The effects of chronic high dose androgen or estrogen treatment on the human prostate. *J Clin Endocrinol Metab* 1996; 81(12):4290-4295.
- Johnson BA, G Marzani-Nissen: Alcohol: Clinical Aspects. B.A. Johnson (ed.), *Addiction Medicine*, Springer Science+Business Media. 2010.
- Jones AL, KJ Simpson: Review article: mechanisms and management of hepatotoxicity in ecstasy (MDMA) and amphetamine intoxications. *Aliment Pharmacol Ther* 1999; 13(2):129-133.

- Jones RT: The pharmacology of cocaine smoking in humans. *NIDA Res Monogr* 1990; 99:30-41.
- Jorgensen E, JT Kaimenyi: The status of periodontal health and oral hygiene of *Miraa* (*catha edulis*) chewers. *East Afr Med J* 1990; 67(8):585-590.
- Kafrouni MI, RA Anders, S Verma: Hepatotoxicity associated with dietary supplements containing anabolic steroids. *Clin Gastroenterol Hepatol* 2007; 5(7):809-812.
- Kaku DA, DH Lowenstein: Emergence of recreational drug abuse as a major risk factor for stroke in young adults. *Ann Intern Med* 1990; 113:821-827.
- Kalsi SS, DM Wood, PI Dargan: The epidemiology and patterns of acute and chronic toxicity associated with recreational ketamine use. *Emerg Health Threats J* 4, 7107-7117, 2011.
- Kam PC, FF Yoong: Gamma-hydroxybutyric acid: an emerging recreational drug. *Anaesthesia* 1998; 53(12):1195-1198.
- Kanayama G, JI Hudson, HGJ Pope: Long-term psychiatric and medical consequences of anabolic-androgenic steroid abuse: a looming public health concern? *Drug Alcohol Depend* 2008; 98(1-2):1-12.
- Kandel DB, M Davies: High school students who use crack and other drugs. *Arch Gen Psychiatry* 1996; 53(1):71-80.
- Kang SY, MF Goldstein, S Deren: Health care utilization and risk behaviors among HIV positive minority drug users. *J Health Care Poor Underserved* 2006; 17(2):265-275.
- Karch SB: *Drug Abuse Handbook*, 2nd ed., CRC Press, Florida, USA. 2006.
- Karch SB, BG Stephens, CH Ho: Methamphetamine-related deaths in San Francisco: demographic, pathologic, and toxicologic profiles. *J Forensic Sci* 1999; 44(2):359-368.
- Karila TAM, JE Karjalainen, MJ Mantysaari, MT Viitasalo, TA Seppala: Anabolic androgenic steroids produce dose-dependant increase in left ventricular mass in power athletes, and this effect is potentiated by concomitant use of growth hormone. *Int J Sports Med* 2003; 24(5):337-343.
- Kassie F, F Darroudi, M Kundi, R Schulte-Hermann, S Knasmuller: Khat (*Catha edulis*) consumption causes genotoxic effects in humans. *Int J Cancer* 2001; 92(3):329-332.
- Kaye BR, M Fainstat: Cerebral vasculitis associated with cocaine abuse. *JAMA* 1987; 258(15):2104-2106.
- Kaye S, S Darke, J Dufloy: Methylenedioxymethamphetamine (MDMA)-related fatalities in Australia: demographics, circumstances, toxicology and major organ pathology. *Drug Alcohol Depend* 2009; 104(3):254-261.
- Kaye S, R McKetin, J Dufloy, S Darke: Methamphetamine and cardiovascular pathology: a review of the evidence. *Addiction* 2007; 102(8):1204-1211.
- Kennedy JG, J Teague, W Rokaw, E Cooney: A medical evaluation of the use of qat in North Yemen. *Soc Sci Med* 1983; 17(12):783-793.
- Kennedy MC, C Lawrence: Anabolic steroid abuse and cardiac death. *Med J Aust* 1993; 158(5):346-348.
- Kinane DF, IG Chestnutt: Smoking and periodontal disease. *Crit Rev Oral Biol Med* 2000; 11(3):356-365.
- Kintz P: Deaths involving buprenorphine: a compendium of French cases. *Forensic Sci Int* 2001; 121(1-2):65-69.
- Klatsky AL: Epidemiology of coronary heart disease--influence of alcohol. *Alcohol Clin Exp Res* 1994; 18(1):88-96.
- Klatsky AL, MA Armstrong, GD Friedman: Red wine, white wine, liquor, beer, and risk for coronary artery disease hospitalization. *Am J Cardiol* 1997; 80(4):416-420.

- Klein RG, S Mannuzza: Hyperactive boys almost grown up. III. Methylphenidate effects on ultimate height. *Arch Gen Psychiatry* 1988; 45(12):1131-1134.
- Kline JN, JD Hirasuna: Pulmonary edema after freebase cocaine smoking--not due to an adulterant. *Chest* 1990; 97(4):1009-1010.
- Klock JC, U Boerner, CE Becker: Coma, hyperthermia and bleeding associated with massive LSD overdose. A report of eight cases. *West J Med* 1974; 120(3):183-188.
- Kopman AF: To the editor: ketamine may interact with barbiturate premedication in producing delayed awakening from anesthesia and significant respiratory depression. *Anesth Analg* 1972; 51(5):793-794.
- Koski A, I Ojanpera, E Vuori: Alcohol and benzodiazepines in fatal poisonings. *Alcohol Clin Exp Res* 2002; 26(7):956-959.
- KPMG: KPMG Economic Consulting. Excessive alcohol consumption in the Netherlands. Trends and social costs. KPMG, Hoofddorp, The Netherlands. www.w3.nigz.nl/docfiles/kostenenbatenkpmp_engels1.pdf. 2001.
- Kraft B, HG Kress: [Cannabinoids and the immune system. Of men, mice and cells]. *Cannabinoide und Immunsystem. Menschen, Mause und Zellen. Schmerz* 2004; 18(3):203-210.
- Kral AH, RN Bluthenthal, RE Booth, JK Watters: HIV seroprevalence among street-recruited injection drug and crack cocaine users in 16 US municipalities. *Am J Public Health* 1998; 88(1):108-113.
- Krendel DA, SM Ditter, MR Frankel, WK Ross: Biopsy-proven cerebral vasculitis associated with cocaine abuse. *Neurology* 1990; 40(7):1092-1094.
- Krieg A, J Scharhag, T Albers, W Kindermann, A Urhausen: Cardiac tissue Doppler in steroid users. *Int J Sports Med* 2007; 28(8):638-643.
- Kriegstein AR, BA Armitage, PY Kim: Heroin inhalation and progressive spongiform leukoencephalopathy. *N Engl J Med* 1997; 336(8):589-590.
- Kummoona R: Surgical reconstruction of the temporomandibular joint for chronic subluxation and dislocation. *Int J Oral Maxillofac Surg* 2001; 30(4):344-348.
- Laborit H: Sodium 4-hydroxybutyrate. *Int J Neuropharmacol* 3, 433-451, 1964.
- Lecci A, F Borsini, L Gragnani, G Volterra, A Meli: Effect of psychotomimetics and some putative anxiolytics on stress-induced hyperthermia. *J Neural Transm Gen Sect* 1991; 83(1-2):67-76.
- Lee CY, H Mohammadi, RA Dixon: Medical and dental implications of cocaine abuse. *J Oral Maxillofac Surg* 1991; 49(3):290-293.
- Leibach WK: Cirrhosis in the alcoholic and its relation to the volume of alcohol abuse. *Ann N Y Acad Sci* 1975; 252:85-105.
- Leone A: Smoking, haemostatic factors, and cardiovascular risk. *Curr Pharm Des* 2007; 13(16):1661-1667.
- Leung SY: Benzodiazepines, opioids and driving: an overview of the experimental research. *Drug Alcohol Rev* 2011; 30(3):281-286.
- Lind J, A Oyefeso, M Pollard, A Baldacchino, H Ghodse: Death rate from use of ecstasy or heroin. *Lancet* 1999; 354(9196):2167.
- Longnecker MP: Alcohol consumption and risk of cancer in humans: an overview. *Alcohol* 1995; 12(2):87-96.
- Longnecker MP, MJ Orza, ME Adams, J Vioque, TC Chalmers: A meta-analysis of alcoholic beverage consumption in relation to risk of colorectal cancer. *Cancer Causes Control* 1990; 1(1):59-68.
- Lopez F, R Silva, L Pestreich, R Muniz: Comparative efficacy of two once daily methylphenidate formulations (Ritalin LA and Concerta) and placebo in children with attention deficit hyperactivity disorder across the school day. *Paediatr Drugs* 2003; 5(8):545-555.

- Lucas GM, M Griswold, KA Gebo, J Keruly, RE Chaisson, RD Moore: Illicit drug use and HIV-1 disease progression: a longitudinal study in the era of highly active antiretroviral therapy. *Am J Epidemiol* 2006; 163:412-420.
- Lucas PB, DL Gardner, OM Wolkowitz, EE Tucker, RW Cowdry: Methylphenidate-induced cardiac arrhythmias. *N Engl J Med* 1986; 315(23):1485.
- Luqman W, TS Danowski: The use of khat (*Catha edulis*) in Yemen. Social and medical observations. *Ann Intern Med* 1976; 85(2):246-249.
- Macigo FG, DL Mwaniki, SW Guthua: The association between oral leukoplakia and use of tobacco, alcohol and khat based on relative risks assessment in Kenya. *Eur J Oral Sci* 1995; 103(5):268-273.
- Makki I: Oral carcinomas and their relationship to khat and shamma abuses. Thesis University of Heidelberg (GDR). 1975.
- Maloney WJ: Significance of cannabis use to dental practice. *N Y State Dent J* 2011; 77(3):36-39.
- Maraj S, VM Figueredo, D Lynn Morris: Cocaine and the heart. *Clin Cardiol* 2010; 33(5):264-269.
- Mark TL, GE Woody, T Juday, HD Kleber: The economic costs of heroin addiction in the United States. *Drug Alcohol Depend* 2001; 61(2):195-206.
- Marker P, A Krogdahl: Plasma cell gingivitis apparently related to the use of khat: report of a case. *Br Dent J* 2002; 192(6):311-313.
- Matick H, D Anderson, J Brumlik: Cerebral vasculitis associated with oral amphetamine overdose. *Arch Neurol* 1983; 40(4):253-254.
- Mattes JA, R Gittelman: Growth of hyperactive children on maintenance regimen of methylphenidate. *Arch Gen Psychiatry* 1983; 40(3):317-321.
- McCabe SE, KJ Brower, BT West, TF Nelson, H Wechsler: Trends in non-medical use of anabolic steroids by U.S. college students: results from four national surveys. *Drug Alcohol Depend* 2007; 90(2-3):243-251.
- Mccann UD, SO Slate, GA Ricaurte: Adverse reactions with 3,4-methylenedioxymethamphetamine (MDMA; 'ecstasy'). *Drug Saf* 1996; 15(2):107-115.
- McKee SA, RJ Applegate, JR Hoyle, MT Sacrinty, MA Kutcher, DC Sane: Cocaine use is associated with an increased risk of stent thrombosis after percutaneous coronary intervention. *Am Heart J* 2007; 154(1):159-164.
- McNutt RA, GS Ferenchick, PC Kirlin, NJ Hamlin: Acute myocardial infarction in a 22-year-old world class weight lifter using anabolic steroids. *Am J Cardiol* 1988; 62(1):164.
- Mehra R, BA Moore, K Crothers, J Tetrault, DA Fiellin: The association between marijuana smoking and lung cancer: a systematic review. *Arch Intern Med* 2006; 166(13):1359-1367.
- Meister KA, EM Whelan, R Kava: The health effects of moderate alcohol intake in humans: an epidemiologic review. *Crit Rev Clin Lab Sci* 2000; 37(3):261-296.
- Melchert RB, AA Welder: Cardiovascular effects of androgenic-anabolic steroids. *Med Sci Sports Exerc* 1995; 27(9):1252-1262.
- Mengel R, M Eigenbrodt, T Schunemann, L Flores-de-Jacoby: Periodontal status of a subject sample of Yemen. *J Clin Periodontol* 1996; 23(5):437-443.
- Menon DK: Successful treatment of anabolic steroid-induced azoospermia with human chorionic gonadotropin and human menopausal gonadotropin. *Fertil Steril* 2003; 79 Suppl 3:1659-1661.
- Menz V, W Grimm, J Hoffmann, B Maisch: Alcohol and rhythm disturbance: the holiday heart syndrome. *Herz* 1996; 21(4):227-231.
- Meyer JS, LF Quenzer: Psychomotor stimulants: Cocaine and the amphetamines. In J.S. Meyer & L.F. Quenzer (Eds.), *Psychopharmacology: Drugs, The Brain, and Behavior* (pp. 275-302). Sunderland, Massachusetts: Sinauer Associates, Inc. Publishers. 2005.

- Milroy CM, JC Clark, AR Forrest: Pathology of deaths associated with 'ecstasy' and 'eve' misuse. *J Clin Pathol* 1996; 49(2):149-153.
- Mitchell-Lewis DA, JA Phelan, RB Kelly, JJ Bradley, IB Lamster: Identifying oral lesions associated with crack cocaine use. *J Am Dent Assoc* 1994; 125(8):1104-8, 1110.
- Mittleman MA, RA Lewis, M Maclure, JB Sherwood, JE Muller: Triggering myocardial infarction by marijuana. *Circulation* 2001; 103(23):2805-2809.
- Mittleman MA, D Mintzer, M Maclure, GH Tofler, JB Sherwood, JE Muller: Triggering of myocardial infarction by cocaine. *Circulation* 1999; 99(21):2737-2741.
- Mixmag: Mixmag Drug Survey 2000-2005 - Evidence includes personal communication from Dr Luke Mitcheson. 2005.
- Modlinski R, KB Fields: The effect of anabolic steroids on the gastrointestinal system, kidneys, and adrenal glands. *Curr Sports Med Rep* 2006; 5(2):104-109.
- Molendijk B, G ter Horst, MB Kasbergen, GJ Truin, J Mulder: [Dental health in drug and alcohol addicts]. *Tandheelkundige gezondheid van drugs- en alcoholverslaafden. Ned Tijdschr Tandheelkd* 1995; 102(8):296-298.
- Molendijk G: [The drug addict and the dentist]. *De verslaafde en de tandarts. Ned Tijdschr Tandheelkd* 1992; 99(2):43-45.
- Moliterno DJ, JE Willard, RA Lange, BH Negus, JD Boehrer, DB Glamann, C Landau, JD Rossen, MD Winniford, LD Hillis: Coronary-artery vasoconstriction induced by cocaine, cigarette smoking, or both. *N Engl J Med* 1994; 330(7):454-459.
- Monfrecola G, G Riccio, C Savarese, G Posteraro, EM Procaccini: The acute effect of smoking on cutaneous microcirculation blood flow in habitual smokers and nonsmokers. *Dermatology* 1998; 197(2):115-118.
- Mooney L, S Glasner-Edwards, RA Rawson, W Ling: Medical effects of methamphetamine use. In: JM Roll, RA Rawson, W Ling W, S Shoptaw (eds) *Methamphetamine addiction: from basic science to treatment*. Guilford, New York, pp 117-142. 2009.
- Moore KA, EM Kilbane, R Jones, GW Kunsman, B Levine, M Smith: Tissue distribution of ketamine in a mixed drug fatality. *J Forensic Sci* 1997; 42(6):1183-1185.
- Morgentaler A: Testosterone and prostate cancer: an historical perspective on a modern myth. *Eur Urol* 2006; 50(5):935-939.
- Morgentaler A: New concepts regarding the relationship of testosterone and prostate cancer. *Future Oncol* 2007; 3(2):123-126.
- Moskal A, T Norat, P Ferrari, E Riboli. Alcohol intake and colorectal cancer risk: a dose-response meta-analysis of published cohort studies. *Int J Cancer* 120, 664-671, 2007.
- Moss AR, JA Hahn, S Perry, ED Charlebois, D Guzman, RA Clark, DR Bangsberg: Adherence to highly active antiretroviral therapy in the homeless population in San Francisco: a prospective study. *Clin Infect Dis* 2004; 39(8):1190-1198.
- Mueller PD, NL Benowitz, KR Olson: Cocaine. *Emerg Med Clin North Am* 1990; 8(3):481-493.
- Mueller PD, WS Korey: Death by 'ecstasy': the serotonin syndrome? *Ann Emerg Med* 1998; 32(3 Pt 1):377-380.
- Muetzelfeldt L, SK Kamboj, H Rees, J Taylor, CJA Morgan, HV Curran: Journey through the K-hole: phenomenological aspects of ketamine use. *Drug Alcohol Depend* 2008; 95(3):219-229.
- Mujili HM, X Bo, L Zhang: The effect of Khat (*Catha edulis*) on acute cerebral infarction. *Neurosci* 10, 219-222, 2005.

- Murphy LL, RM Munoz, BA Adrian, MA Villanua: Function of cannabinoid receptors in the neuroendocrine regulation of hormone secretion. *Neurobiol Dis* 1998; 5(6 Pt B):432-446.
- Mwenda JM, MM Arimi, MC Kyama, DK Langat: Effects of khat (*Catha edulis*) consumption on reproductive functions: a review. *East Afr Med J* 2003; 80(6):318-323.
- Nadel DM, KM Lyons: 'Shotgunning' crack cocaine as a potential cause of retropharyngeal abscess. *Ear Nose Throat J* 1998; 77(1):47-50.
- Nasr AH, ML Khatri: Head and neck squamous cell carcinoma in Hajjah, Yemen. *Saudi Med J* 2000; 21(6):565-568.
- NCETA: National Centre for Education and Training on Addiction Consortium (NCETA). Alcohol and Other Drugs: A Handbook for Health Professionals. Chapter 6 Amphetamines. [http://www.aodgp.gov.au/internet/aodgp/publishing.nsf/Content/handbook/\\$FILE/chap6.pdf](http://www.aodgp.gov.au/internet/aodgp/publishing.nsf/Content/handbook/$FILE/chap6.pdf) . 2004. Australian Government Department of Health and Ageing. 2004.
- NDM: Nationale Drug Monitor (NDM). Jaarbericht 2010. Trimbos, Utrecht. 2011.
- Nencini P, AM Ahmed: Khat consumption: a pharmacological review. *Drug Alcohol Depend* 1989; 23(1):19-29.
- Newton TF, IA Cook, AD Kalechstein, S Duran, F Monroy, W Ling, AF Leuchter: Quantitative EEG abnormalities in recently abstinent methamphetamine dependent individuals. *Clin Neurophysiol* 2003; 114(3):410-415.
- NIAAA: National Institute on Alcohol Abuse and Alcoholism. Medical consequences of alcohol abuse. *Alcohol Res Health* 24, 27-31, 2000.
- NIDA: National Institute on Drug Abuse. *Tob Addiction* 06-4342, pp. 1-8. 2006.
- Nottin S, LD Nguyen, M Terbah, P Obert: Cardiovascular effects of androgenic anabolic steroids in male bodybuilders determined by tissue Doppler imaging. *Am J Cardiol* 2006; 97(6):912-915.
- Novick DM, BL Richman, JM Friedman, JE Friedman, C Fried, JP Wilson, A Townley, MJ Kreek: The medical status of methadone maintenance patients in treatment for 11-18 years. *Drug Alcohol Depend* 1993; 33(3):235-245.
- O'Donnell AE, FG Mappin, TJ Sebo, H Tazelaar: Interstitial pneumonitis associated with 'crack' cocaine abuse. *Chest* 1991; 100(4):1155-1157.
- Oberg M, MS Jaakkola, A Woodward, A Peruga, A Pruss-Ustun: Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet* 2011; 377(9760):139-146.
- Olkola KT, J Ahonen: Midazolam and other benzodiazepines. *Handb Exp Pharmacol* 2008;(182):335-360.
- Oxley JD, AM Cottrell, S Adams, D Gillatt: Ketamine cystitis as a mimic of carcinoma in situ. *Histopathology* 2009; 55(6):705-708.
- Pantelis C, CG Hindler, JC Taylor: Use and abuse of khat (*Catha edulis*): a review of the distribution, pharmacology, side effects and a description of psychosis attributed to khat chewing. *Psychol Med* 1989; 19:657-668.
- Parry J, S Porter, C Scully, S Flint, MG Parry: Mucosal lesions due to oral cocaine use. *Br Dent J* 1996; 180(12):462-464.
- Parssinen M, U Kujala, E Vartiainen, S Sarna, T Seppala: Increased premature mortality of competitive powerlifters suspected to have used anabolic agents. *Int J Sports Med* 2000; 21(3):225-227.
- Parssinen M, T Seppala: Steroid use and long-term health risks in former athletes. *Sports Med* 2002; 32(2):83-94.
- Pechnick RN, JT Ungerleider: Hallucinogens. In J.H. Lowinson, P. Ruiz, R.B. Millman & J.G. Langrod (Eds.), *Substance Abuse: A Comprehensive Textbook* (4th ed.). Baltimore: Williams & Wilkins, pp. 313-323. 2004.

- Pennings EJM, AP Leccese, FA Wolff: Effects of concurrent use of alcohol and cocaine. *Addiction* 2002; 97(7):773-783.
- Perez JAJ, EL Arsura, S Strategos: Methamphetamine-related stroke: four cases. *J Emerg Med* 1999; 17(3):469-471.
- Petitti DB, S Sidney, C Quesenberry, A Bernstein: Stroke and cocaine or amphetamine use. *Epidemiology* 1998; 9(6):596-600.
- Picozzi A, SF Dworkin, JG Leeds, J Nash: J Dental and associated attitudinal aspects of heroin addiction: a pilot study. *J Dent Res* 1972; 51(3):869.
- Pirwitz MJ, JE Willard, C Landau, RA Lange, DB Glamann, DJ Kessler, EH Foerster, E Todd, LD Hillis: Influence of cocaine, ethanol, or their combination on epicardial coronary arterial dimensions in humans. *Arch Intern Med* 1995; 155(11):1186-1191.
- Poon TL, KF Wong, MY Chan, KW Fung, SK Chu, CW Man, MK Yiu, SK Leung: Upper gastrointestinal problems in inhalational ketamine abusers. *J Dig Dis* 2010; 11(2):106-110.
- Pope HGJ, KJ Brower: Anabolic-androgenic steroid-related disorders. In: B Sadock, V Sadock, editors. *Comprehensive textbook of psychiatry*. Ninth Edition. Philadelphia, PA: Lippincott Williams & Wilkins, p. 1419-1431. 2009.
- Pope HGJ, DL Katz: Psychiatric and medical effects of anabolic-androgenic steroid use. A controlled study of 160 athletes. *Arch Gen Psychiatry* 1994; 51(5):375-382.
- Pullan PT, FE Watson, SS Seow, W Rapoport: Methadone-induced hypoadrenalism. *Lancet* 1983; 1(8326 Pt 1):714.
- Quartilho A, HP Mata, MM Ibrahim, TW Vanderah, F Porreca, A Makriyannis, TPJ Malan: Inhibition of inflammatory hyperalgesia by activation of peripheral CB2 cannabinoid receptors. *Anesthesiology* 2003; 99:955-960.
- Quetin AM: *La psilocybine en psychiatrie clinique et experimentale*. Paris: Medical dissertation. 1960.
- Qureshi S, M Tariq, NS Parmar, IA al Meshal: Cytological effects of khat (*Catha edulis*) in somatic and male germ cells of mice. *Drug Chem Toxicol* 1988; 11(2):151-165.
- Ragland AS, Y Ismail, EL Arsura: Myocardial infarction after amphetamine use. *Am Heart J* 1993; 125(1):247-249.
- Rahman MM, I Laher: Structural and functional alteration of blood vessels caused by cigarette smoking: an overview of molecular mechanisms. *Curr Vasc Pharmacol* 2007; 5(4):276-292.
- Rall TW: Hypnotics and sedatives: ethanol. In *Goodman and Gilman's the pharmacological basis of therapeutics*. Chapter 17. McGraw-Hill. 1992.
- Rehm J, B Taylor, J Patra: Volume of alcohol consumption, patterns of drinking and burden of disease in the European region 2002. *Addiction* 2006; 101(8):1086-1095.
- Rehm JT, SJ Bondy, CT Sempos, CV Vuong: Alcohol consumption and coronary heart disease morbidity and mortality. *Am J Epidemiol* 1997; 146(6):495-501.
- Rehm J, R Room, K Graham, M Monteiro, G Gmel, CT Sempos: The relationship of average volume of alcohol consumption and patterns of drinking to burden of disease: an overview. *Addiction* 2003; 98(9):1209-1228.
- Rehm J, B Taylor, S Mohapatra, H Irving, D Baliunas, J Patra, M Roerecke: Alcohol as a risk factor for liver cirrhosis: a systematic review and meta-analysis. *Drug Alcohol Rev* 2010; 29(4):437-445.
- Reich DL, G Silvay: Ketamine: an update on the first twenty-five years of clinical experience. *Can J Anaesth* 1989; 36(2):186-197.

- Reynolds JEF: The extra pharmacopoeia. The pharmaceutical press, London, pp. 1223-1224. 1993.
- Rice EK, NM Isbel, GJ Becker, RC Atkins, LP McMahon: Heroin overdose and myoglobinuric acute renal failure. Clin Nephrol 2000; 54(6):449-454.
- Richards JR, SW Bretz, EB Johnson, SD Turnipseed, BT Brofeldt, RW Derlet: Methamphetamine abuse and emergency department utilization. West J Med 1999; 170(4):198-202.
- Risser D, S Honigschnabl, M Stichenwirth, S Pfufl, D Sebald, A Kaff, G Bauer: Mortality of opiate users in Vienna, Austria. Drug Alcohol Depend 2001; 64(3):251-256.
- RIVM: Nationaal Kompas Volksgezondheid, 2008. Wat zijn de mogelijke gezondheidsgevolgen van roken? Available from: http://www.rivm.nl/vtv/object_document/o1208n19085.html. 2008.
- Roberts JT, DM Essenhigh: Adenocarcinoma of prostate in 40-year-old body-builder. Lancet 1986; 2(8509):742.
- Rogol AD, CE Yesalis: Clinical review 31: Anabolic-androgenic steroids and athletes: what are the issues? J Clin Endocrinol Metab 1992; 74:465-469.
- Room R, T Babor, J Rehm: Alcohol and public health. Lancet 2005; 365(9458):519-530.
- Roth MD, DP Tashkin, R Choi, BD Jamieson, JA Zack, GC Baldwin: Cocaine enhances human immunodeficiency virus replication in a model of severe combined immunodeficient mice implanted with human peripheral blood leukocytes. J Infect Dis 2002; 185(5):701-705.
- Rothrock JF, R Rubenstein, PD Lyden: Ischemic stroke associated with methamphetamine inhalation. Neurology 1988; 38(4):589-592.
- Ruiz P, E Strain: Lowinson and Ruiz's Substance Abuse. A Comprehensive Textbook. Section VII, Management of associated medical conditions, pp. 850. 2011.
- Rumbaugh CL, RT Bergeron, RL Scanlan, JS Teal, HD Segall, HC Fang, R McCormick: Cerebral vascular changes secondary to amphetamine abuse in the experimental animal. Radiology 1971; 101(2):345-351.
- Rumbaugh CL, HC Fang, RE Higgins, RT Bergeron, HD Segall, JS Teal: Cerebral microvascular injury in experimental drug abuse. Invest Radiol 1976; 11(4):282-294.
- Safdar K, ER Schiff: Alcohol and hepatitis C. Semin Liver Dis 2004; 24:305-315.
- Sainsbury D: Drug addiction and dental care. N Z Dent J 1999; 95(420):58-61.
- Sakkour A, T Wang, D Tashkin: A 56-year-old woman with COPD and multiple pulmonary nodules. Chest 2008; 133(2):566-569.
- Sanchez-Osorio M, A Duarte-Rojo, B Martinez-Benitez, A Torre, M Uribe: Anabolic-androgenic steroids and liver injury. Liver Int 2008; 28(2):278-282.
- Sano R, T Hasuike, M Nakano, Y Kominato, H Itoh: A fatal case of myocardial damage due to misuse of the 'designer drug' MDMA. Leg Med (Tokyo) 2009; 11(6):294-297.
- Santora LJ, J Marin, J Vangrow, C Minegar, M Robinson, J Mora, G Friede: Coronary calcification in body builders using anabolic steroids. Prev Cardiol 2006; 9(4):198-201.
- Sarnquist F, CPJ Larson: Drug-induced heat stroke. Anesthesiology 1973; 39(3):348-350.
- Sasco AJ, RM Merrill, I Dari, V Haim-Luzon, F Carriot, CI Cann, M Bartal: A case-control study of lung cancer in Casablanca, Morocco. Cancer Causes Control 2002; 13(7):609-616.
- Saxer UP, C Walter, MM Bornstein, K Klingler, CA Ramseier: [Impact of tobacco use on the periodontium--an update. Part 2: Clinical and radiographic

- changes in the periodontium and effects on periodontal and implant therapy]
- Einfluss des Tabakkonsums auf das Parodont--ein Update. II. Teil 2: Klinische und radiologische Veränderungen des Parodonts sowie Folgen auf die Parodontaltherapie und orale Implantologie. Schweiz Monatsschr Zahnmed 2007; 117(2):153-169.
- Scheutz F: Five-year evaluation of a dental care delivery system for drug addicts in Denmark. Community Dent Oral Epidemiol 1984; 12(1):29-34.
- Schoser BG, C Groden: Subacute onset of oculogyric crises and generalized dystonia following intranasal administration of heroin. Addiction 1999; 94(3):431-434.
- Schuckit MA: Drug and Alcohol Abuse: A Clinical Guide to Diagnosis and Treatment (Critical Issues in Psychiatry). Chapter 2 Depressants. Kluwer Academic Plenum Publ. 5th Ed. 2000.
- Serfaty M, G Masterton: Fatal poisonings attributed to benzodiazepines in Britain during the 1980s. Br J Psychiatry 1993; 163:386-393.
- Shahani R, C Streutker, B Dickson, RJ Stewart: Ketamine-associated ulcerative cystitis: a new clinical entity. Urology 2007; 69(5):810-812.
- Shaner JW, N Kimmes, T Saini, P Edwards: 'Meth mouth': rampant caries in methamphetamine abusers. AIDS Patient Care STDS 2006; 20:146-150.
- Shaner JW: Caries associated with methamphetamine abuse. J Mich Dent Assoc 2002; 84(9):42-47.
- Shaw HEJ, JG Lawson, RD Stulting: Amaurosis fugax and retinal vasculitis associated with methamphetamine inhalation. J Clin Neuroophthalmol 1985; 5(3):169-176.
- Sherrill DL, M Krzyzanowski, JW Bloom, MD Lebowitz: Respiratory effects of non-tobacco cigarettes: a longitudinal study in general population. Int J Epidemiol 1991; 20(1):132-137.
- Shifren JL: The role of androgens in female sexual dysfunction. Mayo Clin Proc 2004; 79(4 Suppl):S19-S24.
- Sidney S, CPJ Quesenberry, GD Friedman, IS Tekawa: Marijuana use and cancer incidence (California, United States). Cancer Causes Control 1997; 8(5):722-728.
- Sidney S: Cardiovascular consequences of marijuana use. J Clin Pharmacol 2002; 42(11 Suppl):64S-70S.
- Siegel RK: Phencyclidine and ketamine intoxication: a study of four populations of recreational users. NIDA Res Monogr 1978;(21):119-147.
- Singh AB, S Hsia, P Alaupovic, I Sinha-Hikim, L Woodhouse, TA Buchanan, R Shen, R Bross, N Berman, S Bhasin: The effects of varying doses of T on insulin sensitivity, plasma lipids, apolipoproteins, and C-reactive protein in healthy young men. J Clin Endocrinol Metab 2002; 87(1):136-143.
- Skeie I, M Brekke, M Lindbaek, H Waal: Somatic health among heroin addicts before and during opioid maintenance treatment: a retrospective cohort study. BMC Public Health 2008; 8:43.
- Smith GT, PL McClaughry, J Purkey, W Thompson: Crack cocaine mimicking pulmonary embolism on pulmonary ventilation/perfusion lung scan. A case report. Clin Nucl Med 1995; 20(1):65-68.
- Smith HJ, AH Roche, MF Jausch, PB Herdson: Cardiomyopathy associated with amphetamine administration. Am Heart J 1976; 91(6):792-797.
- Smith HS: Ketamine-induced urologic insult (KIUI). Pain Physician 2010; 13(6):E343-E346.
- Smith-Warner SA, D Spiegelman, SS Yaun, PA van den Brandt, AR Folsom, RA Goldbohm, S Graham, L Holmberg, GR Howe, JR Marshall, AB Miller, JD Potter, FE Speizer, WC Willett, A Wolk, DJ Hunter: Alcohol and breast

- cancer in women: a pooled analysis of cohort studies. *JAMA* 1998; 279(7):535-540.
- Socas L, M Zumbado, O Perez-Luzardo, A Ramos, C Perez, JR Hernandez, LD Boada: Hepatocellular adenomas associated with anabolic androgenic steroid abuse in bodybuilders: a report of two cases and a review of the literature. *Br J Sports Med* 2005; 39(5):e27.
- Sorensen TI, M Orholm, KD Bentsen, G Hoybye, K Eghoje, P Christoffersen: Prospective evaluation of alcohol abuse and alcoholic liver injury in men as predictors of development of cirrhosis. *Lancet* 1984; 2(8397):241-244.
- Soufi HE, M Kameswaran, T Malatani: Khat and oral cancer. *J Laryngol Otol* 1991; 105(8):643-645.
- Sparacino CM, PA Hyldborg, TJ Hughes: Chemical and biological analysis of marijuana smoke condensate. *NIDA Res Monogr* 1990; 99:121-140.
- Sporer KA: The serotonin syndrome. Implicated drugs, pathophysiology and management. *Drug Saf* 1995; 13(2):94-104.
- Sporer KA: Acute heroin overdose. *Ann Intern Med* 1999; 130(7):584-590.
- Sporer KA, RL Chin, JE Dyer, R Lamb: Gamma-hydroxybutyrate serum levels and clinical syndrome after severe overdose. *Ann Emerg Med* 42, 3-8, 2003.
- Story A, G Bothamley, A Hayward: Crack cocaine and infectious tuberculosis. *Emerg Infect Dis* 2008; 14(9):1466-1469.
- Strassman RJ, CR Qualls, LM Berg: Differential tolerance to biological and subjective effects of four closely spaced doses of N,N-dimethyltryptamine in humans. *Biol Psychiatry* 1996; 39(9):784-795.
- Sullivan ML, CM Martinez, P Gennis, EJ Gallagher: The cardiac toxicity of anabolic steroids. *Prog Cardiovasc Dis* 1998; 41(1):1-15.
- Sumnall HR, K Woolfall, S Edwards, JC Cole, CM Beynon: Use, function, and subjective experiences of gamma-hydroxybutyrate (GHB). *Drug Alcohol Depend* 92, 286-290, 2008.
- Swan GE, CN Lessov-Schlaggar: The effects of tobacco smoke and nicotine on cognition and the brain. *Neuropsychol Rev* 2007; 17(3):259-273.
- Takahara J, S Yunoki, W Yakushiji, Y Yamane: Stimulatory effects of GHB on growth hormone and prolactin release in humans. *J Clin Endocrinol Metab* 44, 1014-1017, 1997.
- Tan WC, C Lo, A Jong, L Xing, MJ Fitzgerald, WM Vollmer, SA Buist, DD Sin: Marijuana and chronic obstructive lung disease: a population-based study. *CMAJ* 2009; 180(8):814-820.
- Tardiff K, E Gross, J Wu, M Stajic, R Millman: Analysis of cocaine-positive fatalities. *J Forensic Sci* 1989; 34(1):53-63.
- Tariq M, S Qureshi, AM Ageel, IA al Meshal: The induction of dominant lethal mutations upon chronic administration of khat (*Catha edulis*) in albino mice. *Toxicol Lett* 1990; 50(2-3):349-353.
- Tashkin DP: Airway effects of marijuana, cocaine, and other inhaled illicit agents. *Curr Opin Pulm Med* 2001; 7(2):43-61.
- Tashkin DP: Is frequent marijuana smoking harmful to health? *West J Med* 1993; 158(6):635-637.
- Tashkin DP: Smoked marijuana as a cause of lung injury. *Monaldi Arch Chest Dis* 2005; 63(2):93-100.
- Tashkin DP, AH Coulson, VA Clark, M Simmons, LB Bourque, S Duann, GH Spivey, H Gong: Respiratory symptoms and lung function in habitual heavy smokers of marijuana alone, smokers of marijuana and tobacco, smokers of tobacco alone, and nonsmokers. *Am Rev Respir Dis* 1987; 135(1):209-216.

- Tashkin DP, EC Kleerup, SN Koyal, JA Marques, MD Goldman: Acute effects of inhaled and i.v. cocaine on airway dynamics. *Chest* 1996; 110(4):904-910.
- Tashkin DP, BJ Shapiro, IM Frank: Acute pulmonary physiologic effects of smoked marijuana and oral Δ^9 -tetrahydrocannabinol in healthy young men. *N Engl J Med* 1973; 289(7):336-341.
- Tashkin DP, BJ Shapiro, IM Frank: Acute effects of smoked marijuana and oral Δ^9 -tetrahydrocannabinol on specific airway conductance in asthmatic subjects. *Am Rev Respir Dis* 1974; 109(4):420-428.
- Tashkin DP, BJ Shapiro, YE Lee, CE Harper: Effects of smoked marijuana in experimentally induced asthma. *Am Rev Respir Dis* 1975; 112:377-386.
- Tashkin DP, MS Simmons, DL Sherrill, AH Coulson: Heavy habitual marijuana smoking does not cause an accelerated decline in FEV1 with age. *Am J Respir Crit Care Med* 1997; 155(1):141-148.
- Taylor DR, R Poulton, TE Moffitt, P Ramankutty, MR Sears: The respiratory effects of cannabis dependence in young adults. *Addiction* 2000; 95(11):1669-1677.
- Taylor DR, DM Fergusson, BJ Milne, LJ Horwood, TE Moffitt, MR Sears, R Poulton: A longitudinal study of the effects of tobacco and cannabis exposure on lung function in young adults. *Addiction* 2002; 97(8):1055-1061.
- Taylor FM: Marijuana as a potential respiratory tract carcinogen: a retrospective analysis of a community hospital population. *South Med J* 1988; 81(10):1213-1216.
- Ter Horst G, B Molendijk, MB Kasbergen: Dental health behavior of drug addicts. *J Dent Res* 237, 545-551, 1999.
- Tetrault JM, K Crothers, BA Moore, R Mehra, J Concato, DA Fiellin: Effects of marijuana smoking on pulmonary function and respiratory complications: a systematic review. *Arch Intern Med* 2007; 167(3):221-228.
- Thadani PV: NIDA conference report on cardiopulmonary complications of 'crack' cocaine use. Clinical manifestations and pathophysiology. *Chest* 1996; 110(4):1072-1076.
- Thiblin I, A Petersson: Pharmacoepidemiology of anabolic androgenic steroids: a review. *Fundam Clin Pharmacol* 2005; 19(1):27-44.
- Thompson PD, EM Cullinane, SP Sady, C Chenevert, AL Saritelli, MA Sady, PN Herbert: Contrasting effects of testosterone and stanozolol on serum lipoprotein levels. *JAMA* 1989; 261(8):1165-1168.
- Thomson WM, R Poulton, JM Broadbent: Cannabis smoking and periodontal disease among young adults. *JAMA* 299, 525-531, 2008.
- Thun MJ, R Peto, AD Lopez, JH Monaco, SJ Henley, CWJ Heath, R Doll: Alcohol consumption and mortality among middle-aged and elderly U.S. adults. *N Engl J Med* 1997; 337(24):1705-1714.
- Togna GI, AR Togna, M Graziani, M Franconi: Testosterone and cocaine: vascular toxicity of their concomitant abuse. *Thromb Res* 2003; 109(4):195-201.
- Tracqui A, P Kintz, B Ludes: Buprenorphine-related deaths among drug addicts in France: a report on 20 fatalities. *J Anal Toxicol* 1998; 22(6):430-434.
- Treadwell SD, TG Robinson: Cocaine use and stroke. *Postgrad Med J* 2007; 83(980):389-394.
- Trimbos: Nationale Drugsmonitor. Jaarbericht 2004. Trimbos institute, Utrecht, The Netherlands. 2004.
- Trugman JM: Cerebral arteritis and oral methylphenidate. *Lancet* 1988; 1(8585):584-585.

- Turek PJ, RH Williams, JH Gilbaugh, LI Lipshultz: The reversibility of anabolic steroid-induced azoospermia. *J Urol* 1995; 153(5):1628-1630.
- Turnipseed SD, JR Richards, JD Kirk, DB Diercks, EA Amsterdam: Frequency of acute coronary syndrome in patients presenting to the emergency department with chest pain after methamphetamine use. *J Emerg Med* 2003; 24(4):369-373.
- Uhde TW, Tancer ME, Shea CA: Sexual dysfunction related to alprazolam treatment of social phobia. *Am J Psychiatry* 1988; 145(4):531-532.
- UKDH: UK Department of Health. Sensible Drinking. The Report of an Inter-Departmental Working Group. Wetherby. 1995.
- USDHHS: U.S.Department of Health and Human Services (USDHHS). The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. 2006.
- Vadivelu N, RL Hines: Buprenorphine: a unique opioid with broad clinical applications. *J Opioid Manag* 2007; 3(1):49-58.
- Van Amsterdam JGC, A Opperhuizen, W van den Brink: Harm potential of magic mushroom use: A review. *Regul Toxicol Pharmacol* 2011.
- Van Baal PH, R Heijink, RT Hoogenveen, JJ Polder. Zorgkosten van ongezond gedrag. RIVM RIVM-rapportnummer: 270751015. <http://www.rivm.nl/bibliotheek/rapporten/270751015.pdf>. 2006.
- Van Baal PH, JJ Polder, GA de Wit, RT Hoogenveen, TL Feenstra, HC Boshuizen, PM Engelfriet, WB Brouwer: Lifetime medical costs of obesity: prevention no cure for increasing health expenditure. . 2008 5(2):e29. *PLoS Med* 5, e29, 2008.
- Van Breda E, HA Keizer, H Kuipers, BHR Wolffenbuttel: Androgenic anabolic steroid use and severe hypothalamic-pituitary dysfunction: a case study. *Int J Sports Med* 2003; 24(3):195-196.
- Van Hoozen BE, CE Cross: Marijuana. Respiratory tract effects. *Clin Rev Allergy Immunol* 1997; 15(3):243-269.
- van Laar MW, AAN Cruys, JEE Verdurmen, MMJ van Ooyen, RF Meyer. Nationale Drug Monitor. Nationale Drug Monitor: Jaarbericht 2007. Utrecht.: Trimboos-instituut. 2008.
- Van Sassenbroek DK, PA Calle, FM Rousseau, AG Verstraete, FM Belpaire, KG Monsieurs, R Haentjes, J Allonsius, J van Brantegem, W Haenen, WA van Buylaert. Medical problems related to recreational drug use at nocturnal dance parties. *Eur J Emerg Med* 10, 302-308, 2003.
- Van Wijk AJ, GHW Verrips, JM Kieffer, J Oudejans, G Molendijk. Mondgezondheidgerelateerde levenskwaliteit van verslaafden. *NedTijdschr Tandheelkd* 118, 219-221, 2011.
- Velazquez I, BP Alter: Androgens and liver tumors: Fanconi's anemia and non-Fanconi's conditions. *Am J Hematol* 2004; 77(3):257-267.
- Vigano D, M Grazia Cascio, T Rubino, F Fezza, A Vaccani, V Di Marzo, D Parolaro: Chronic morphine modulates the contents of the endocannabinoid, 2-arachidonoyl glycerol, in rat brain. *Neuropsychopharmacology* 2003; 28(6):1160-1167.
- Vilensky W: Illicit and licit drugs causing perforation of the nasal septum. *J Forensic Sci* 1982; 27(4):958-962.
- Vittinghoff E, NA Hessel, P Bacchetti, RE Fusaro, SD Holmberg, SP Buchbinder: Cofactors for HIV disease progression in a cohort of homosexual and bisexual men. *J Acquir Immune Defic Syndr* 2001; 27(3):308-314.
- Walter FG, TA Bey, DS Ruschke, NL Benowitz: Marijuana and hyperthermia. *J Toxicol Clin Toxicol* 1996; 34(2):217-221.
- Wang GJ, ND Volkow, L Chang, E Miller, M Sedler, R Hitzemann, W Zhu, J Logan, Y Ma, JS Fowler: Partial recovery of brain metabolism in

- methamphetamine abusers after protracted abstinence. *Am J Psychiatry* 2004; 161(2):242-248.
- Ward J, S Darke, W Hall, R Mattick: Methadone maintenance and the human immunodeficiency virus: current issues in treatment and research. *Br J Addict* 1992; 87(3):447-453.
- Ward J, W Hall, RP Mattick: Role of maintenance treatment in opioid dependence. *Lancet* 1999; 353(9148):221-226.
- Webber MP, EE Schoenbaum, MN Gourevitch, D Buono, RS Klein: A prospective study of HIV disease progression in female and male drug users. *AIDS* 1999; 13(2):257-262.
- Weber JE, CR Chudnofsky, M Boczar, EW Boyer, MD Wilkerson, JE Hollander: Cocaine-associated chest pain: how common is myocardial infarction? *Acad Emerg Med* 2000; 7(8):873-877.
- Weiner AL, L Vieira, CA Mckay, MJ Bayer: Ketamine abusers presenting to the emergency department: a case series. *J Emerg Med* 2000; 18:447-451.
- Weiner WJ, PA Nausieda, HL Klawans: Methylphenidate-induced chorea: case report and pharmacologic implications. *Neurology* 1978; 28:1041-1044.
- Welder AA, JW Robertson, RB Melchert: Toxic effects of anabolic-androgenic steroids in primary rat hepatic cell cultures. *J Pharmacol Toxicol Methods* 1995; 33(4):187-195.
- Welling TH, DM Williams, JC Stanley: Excessive oral amphetamine use as a possible cause of renal and splanchnic arterial aneurysms: a report of two cases. *J Vasc Surg* 1998; 28(4):727-731.
- Wemyss-Holden SA, FC Hamdy, KJ Hastie: Steroid abuse in athletes, prostatic enlargement and bladder outflow obstruction--is there a relationship? *Br J Urol* 1994; 74(4):476-478.
- Westover AN, S McBride, RW Haley: Stroke in young adults who abuse amphetamines or cocaine: a population-based study of hospitalized patients. *Arch Gen Psychiatry* 2007; 64(4):495-502.
- Westover AN, PA Nakonezny, RW Haley: Acute myocardial infarction in young adults who abuse amphetamines. *Drug Alcohol Depend* 2008; 96(1-2):49-56.
- White JM, RJ Irvine: Mechanisms of fatal opioid overdose. *Addiction* 1999; 94(7):961-972.
- White SR, CM Yadao: Characterization of methylphenidate exposures reported to a regional poison control center. *Arch Pediatr Adolesc Med* 2000; 154(12):1199-1203.
- WHO: Review of the pharmacology of khat. Report of a WHO advisory group. *Bull Narc* 1980; 32(3):83-93.
- WHO. Global Status Report on Alcohol (summary). World Health Organisation (WHO).
http://www.who.int/substance_abuse/who_ncd_msd_2001_2.pdf. 2001.
- WHO. World Health Organization. The World Health Report 2003: Shaping the Future. 2003. Geneva, WHO. 2003.
- WHO. WHO Expert Committee on Problems Related to Alcohol Consumption. Second report 2006. Geneva, Switzerland. WHO technical report series; no. 944.
http://www.who.int/substance_abuse/expert_committee_alcohol_trs944.pdf. 2006.
- WHO. Atlas on substance abuse: Resources for the prevention and treatment of substance use disorders. WHO.
http://www.who.int/substance_abuse/publications/Media/en/. 2010.
- WHO. mhGAP newsletter - Mental Health Gap Action Programme. January 2011
http://www.who.int/mental_health/en/. 2011.

- WHO IARC. Tobacco Smoking and Involuntary Smoking, IARC Press, Geneva. 2004.
- Wijetunga M, R Bhan, J Lindsay, S Karch: Acute coronary syndrome and crystal methamphetamine use: a case series. *Hawaii Med J* 2004; 63(1):8-13, 25.
- Wilford JA, NL Day, MD Cornelius. Tobacco use during pregnancy: epidemiology and effects on offspring. In: Miller, M.W. (Ed), *Brain development : normal processes and the effects of alcohol and nicotine*, Oxford, New York, pp. 315-27. 2006.
- Williams H, L Dratcu, R Taylor, M Roberts, A Oyefeso: 'Saturday night fever': ecstasy related problems in a London accident and emergency department. *J Accid Emerg Med* 1998; 15(5):322-326.
- Wilson DJ. Androgens. *Goodman en Gilman's de pharmacological basis van therapeutics*. McGraw-Hill Inc. 8th volume , 1413-1462. 1992.
- Wilson JD: Androgen abuse by athletes. *Endocr Rev* 1988; 9(2):181-199.
- Winecoff AP, RJ Hariman, JJ Grawe, Y Wang, JL Bauman: Reversal of the electrocardiographic effects of cocaine by lidocaine. Part 1. Comparison with sodium bicarbonate and quinidine. *Pharmacotherapy* 1994; 14(6):698-703.
- Withington DE, JA Patrick, F Reynolds: Histamine release by morphine and diamorphine in man. *Anaesthesia* 1993; 48(1):26-29.
- Witton K: On the use of parenteral methylphenidate: a follow-up report. *Am J Psychiatry* 1964; 121:267-268.
- Wolters EC, GK van Wijngaarden, FC Stam, H Rengelink, RJ Lousberg, ME Schipper, B Verbeeten: Leucoencephalopathy after inhaling 'heroin' pyrolysate. *Lancet* 1982; 2(8310):1233-1237.
- Woods BT, GJ Strewler: Hemiparesis occurring six hours after intravenous heroin injection. *Neurology* 1972; 22(8):863-866.
- Wu TC, DP Tashkin, B Djahed, JE Rose: Pulmonary hazards of smoking marijuana as compared with tobacco. *N Engl J Med* 1988; 318:347-351.
- Zelger JL, HX Schorno, EA Carlini: Behavioural effects of cathinone, an amine obtained from *Catha edulis* Forsk.: comparisons with amphetamine, norpseudoephedrine, apomorphine and nomifensine. *Bull Narc* 1980; 32(3):67-81.
- Zhang ZF, H Morgenstern, MR Spitz, DP Tashkin, GP Yu, JR Marshall, TC Hsu, SP Schantz: Marijuana use and increased risk of squamous cell carcinoma of the head and neck. *Cancer Epidemiol Biomarkers Prev* 1999; 8(12):1071-1078.
- Zhou W, PH Lin, RL Bush, L Nguyen, AB Lumsden: Acute arterial thrombosis associated with cocaine abuse. *J Vasc Surg* 2004; 40(2):291-295.
- Zou S, M TepperM, A Giulivi: Current status of hepatitis C in Canada. *Can J Public Health* 2000; 91 Suppl 1:S10-S16.



Published by:

**National Institute for Public Health
and the Environment**

P.O. Box 1 | 3720 BA Bilthoven

The Netherlands

www.rivm.com